arXiv:0810.1840v1 [math.RT] 10 Oct 2008

A BASIC SET FOR THE ALTERNATING GROUP

OLIVIER BRUNAT AND JEAN-BAPTISTE GRAMAIN

Abstract. This article concerns the p-basic set existence problem in the representation theory of finite groups. We show that, for any odd prime p, the alternating group $\mathfrak{A}_n$ has a p-basic set. More precisely, we prove that the symmetric group $\mathfrak{S}_n$ has a p-basic set with some additional properties, allowing us to deduce a p-basic set for $\mathfrak{A}_n$. Our main tool is the generalized perfect isometries introduced by Külshammer, Olsson and Robinson. As a consequence we obtain some results on the decomposition number of $\mathfrak{A}_n$.

1. Introduction

Let G be a finite group and p be a prime. We are interested in the representation theory of G over a field of characteristic p. In the framework of Brauer's modular representation theory, we can construct a decomposition map $d_p : \mathcal{R}_0(G) \to \mathcal{R}_p(G)$ from the Grothendieck group of finite-dimensional representations of G in characteristic 0 to the analogous Grothendieck group in characteristic p. A fundamental result of Brauer's shows that this map is surjective. This motivates the following definition. A set B of irreducible representations of G in characteristic 0 is called a *p-basic set* of G if the images of the classes of the elements in B under d_p form a basis of $\mathcal{R}_p(G)$. Note that it is not clear if p-basic sets always exist. The p-basic sets are powerful tools to compute the p-decomposition matrix of G. For example, Hiss in [9] and Geck in [3] used them to compute the p-decomposition matrices of $G_2(q)$ and $SU_3(q^2)$ respectively, when p is not the defining characteristic. It is an open question however if such p-basic sets exist in general. The answer is known to be positive in the following situations: for p-soluble groups (this is a direct consequence of the result of Fong-Swan in [2, X.2.1]), for finite groups of Lie type in the non-defining characteristics under some additional hypotheses (cf [6], [4] and [5]), for the finite general linear groups $\mathrm{GL}_2(q)$, $\mathrm{GL}_3(q)$ and $\mathrm{GL}_4(q)$ in the defining characteristic (cf [1]), and also for the symmetric group $\mathfrak{S}_n$ (cf [11, 6.3.60]).

This question is still open for the alternating group $\mathfrak{A}_n$. In the present work, we prove that p-basic sets do exist for $\mathfrak{A}_n$, whenever p is an odd prime. For every subset B of irreducible characters of $\mathfrak{S}_n$, we denote by $B_{\mathfrak{A}_n}$ the set of all irreducible constituents of any of the $\mathrm{Res}_{\mathfrak{A}_n}^{\mathfrak{S}_n}(\chi)$'s ($\chi \in B$). We say that $B_{\mathfrak{A}_n}$ is the *restriction* of B to $\mathfrak{A}_n$. Our approach is to find a p-basic set B of $\mathfrak{S}_n$ which restricts to a p-basic set of $\mathfrak{A}_n$. If B is a p-basic set of $\mathfrak{S}_n$, the set $B_{\mathfrak{A}_n}$ is in general not a p-basic set of $\mathfrak{A}_n$. For example, the p-basic set of $\mathfrak{S}_n$ described in [11, 6.3.60] does not restrict to a p-basic set of $\mathfrak{A}_n$.

In [4, §4], Geck gives conditions on B ensuring that $B_{\mathfrak{A}_n}$ is a p-basic set. More precisely, he proves that, for any finite group G with a normal subgroup H such that G/H is abelian, and for any prime p not dividing $|G/H|$, if there is a $\mathcal{R}_0(G/H)$-stable p-basic set B of G such that the set $d_p(B_H)$ is a system of generators of

$\mathcal{R}_p(H)$, then B_H is a p-basic set of H. To prove that $\mathfrak{A}_n$ has a p-basic set, we will construct a p-basic set of $\mathfrak{S}_n$ which satisfies Geck's conditions.

In order to state our main result, we first recall some results and notations. The conjugacy classes and irreducible characters of $\mathfrak{S}_n$ are canonically labelled by the partitions of n (cf [11, 2.1.11]). For λ a partition of n (written $\lambda \vdash n$), we write χ_λ the corresponding irreducible character of $\mathfrak{S}_n$. We denote by ε the sign character of $\mathfrak{S}_n$. This is the linear character of $\mathfrak{S}_n$ which extends the non-trivial character of $\mathrm{Irr}(\mathfrak{S}_n/\mathfrak{A}_n)$. Moreover if $\chi_\lambda \in \mathrm{Irr}(\mathfrak{S}_n)$, then $\varepsilon\chi_\lambda = \chi_{\lambda^*}$, where λ^* denotes the conjugate partition of λ. If $\lambda = \lambda^*$, then we say that λ is *self-conjugate*.

Now let p be a prime. A partition λ of n is said to be *p-regular* if it has no parts divisible by p (because λ then labels a conjugacy class of p-regular elements of $\mathfrak{S}_n$). Note that, sometimes, one also calls *p-regular* any partition of n with no part repeated more than $p-1$ times (cf [11, 6.1]). With this terminology, the set of p-regular partitions of n labels the p-basic set of $\mathfrak{S}_n$ in [11, 6.3.60]. However, as we mentionned above, this basic set does not restrict to a basic set of $\mathfrak{A}_n$, and we will never in this paper use p-regular in this sense.

For $\lambda \vdash n$, we denote by $\gamma(\lambda)$ its p-core and by $\alpha_\lambda = (\lambda^{(1)}, \ldots, \lambda^{(p)})$ its p-quotient. In the following, the i-th part $\lambda^{(i)}$ of the p-quotient α_λ will also be denoted by α_λ^i. Note that the p-quotient of λ is not uniquely defined, but depends on a convention (or a choice of origin). This is described more precisely in §3.1. However, one can show (cf Lemma 3.1 and Convention 3.2) that, for a certain choice of origin, the p-quotient of λ^* is given by $\alpha_{\lambda^*} = (\lambda^{(p)^*}, \ldots, \lambda^{(1)^*})$ (and this independantly on the choice of λ, or even of n).

Finally, to each self-conjugate partition $\lambda = (\lambda_1, \ldots, \lambda_k)$ of n, we associate the partition $\overline{\lambda}$ of n given by

$$\overline{\lambda} = (2\lambda_1 - 1, 2\lambda_2 - 3, \ldots, 2\lambda_k - (2k-1)).$$

Note that the parts of $\overline{\lambda}$ are given by the lengths of the *diagonal hooks* in the Young diagram of λ (i.e. those hooks whose top left corner lies on the diagonal). Our main result is:

Theorem 1.1. *We keep the notation as above. Let p be an odd prime and let $n \in \mathbb{N}$. We set*

$$\Lambda_\emptyset = \{\lambda \vdash n \mid \alpha_\lambda^{(p+1)/2} = \emptyset\}.$$

Then the subset $\mathcal{B}_\emptyset = \{\chi_\lambda \mid \lambda \in \Lambda_\emptyset\}$ is a p-basic set of the symmetric group $\mathfrak{S}_n$. Furthermore, $\mathcal{B}_\emptyset$ satisfies the following properties

1. *If $\chi_\lambda \in \mathcal{B}_\emptyset$, then $\chi_{\lambda^*} \in \mathcal{B}_\emptyset$.*
2. *If $\lambda = \lambda^*$, then $\chi_\lambda \in \mathcal{B}_\emptyset$ if and only if the partition $\overline{\lambda}$ is p-regular.*

Note that the additional properties (1) and (2) are direct consequences of Convention 3.2 (cf Lemma 3.4). The hardest part is to prove that $\mathcal{B}_\emptyset$ is indeed a p-basic set. As a consequence we will prove that $\mathcal{B}_{\emptyset,\mathfrak{A}_n}$, the restriction of the p-basic set $\mathcal{B}_\emptyset$ obtained in the above theorem, is a p-basic set of $\mathfrak{A}_n$ (see Theorem 5.2).

In order to do this, we will need a more general concept of basic sets, which goes as follows. Let $\mathcal{C}$ denote a union of conjugacy classes of G. For every class function φ on G, we define a class function $\varphi^{\mathcal{C}}$ by letting $\varphi^{\mathcal{C}}(g) = \varphi(g)$ if $g \in \mathcal{C}$, and $\varphi^{\mathcal{C}}(g) = 0$ otherwise. Let b be a subset of $\mathrm{Irr}(G)$. A *$\mathcal{C}$-basic set* of b is a subset $B \subseteq b$, such that the family $B^{\mathcal{C}} = \{\chi^{\mathcal{C}} | \chi \in B\}$ is a $\mathbb{Z}$-basis of the $\mathbb{Z}$-module generated by $b^{\mathcal{C}} = \{\chi^{\mathcal{C}} | \chi \in b\}$. If b is the whole of $\mathrm{Irr}(G)$, then a $\mathcal{C}$-basic set of b is just called $\mathcal{C}$-basic set of G (or for G). Moreover, we note that if p is a prime, and if we take $\mathcal{C}$ to be the set of p-regular elements of G, then we write $\mathcal{C} = p$-reg, and $\mathcal{C}$-basic sets are the same as p-basic sets as defined before.

We now present the notions of *generalized blocks* and *generalized perfect isometry* introduced by Külshammer, Olsson and Robinson in [12]. Let $\mathcal{C}$ be a union of conjugacy classes of a finite group G. For any complex-valued class functions α, β of G, we let

$$\langle\, \alpha, \beta \,\rangle_{\mathcal{C}} = \frac{1}{|G|} \sum_{g \in \mathcal{C}} \alpha(g)\overline{\beta(g)}.$$

If $\langle\, \alpha, \beta \,\rangle_{\mathcal{C}} = 0$, then α and β are said to be *orthogonal across* $\mathcal{C}$. Note that, if χ, $\varphi \in \operatorname{Irr}(G)$, then $\langle\, \chi, \varphi \,\rangle_{\mathcal{C}} = \langle\, \chi^{\mathcal{C}}, \varphi \,\rangle_G = \langle\, \chi^{\mathcal{C}}, \varphi^{\mathcal{C}} \,\rangle_G$, where $\chi^{\mathcal{C}}$ and $\varphi^{\mathcal{C}}$ are the class functions we defined before, and $\langle\ ,\ \rangle_G$ is the ordinary scalar product on class functions of G.

We define the *$\mathcal{C}$-blocks* of G to be the minimal subsets of $\operatorname{Irr}(G)$ subject to being orthogonal across $\mathcal{C}$. In particular, it is well-known that, if $\mathcal{C}$ is the set of p-regular elements of G, then the $\mathcal{C}$-blocks are just the p-blocks of modular representation theory. Note also that the $\mathcal{C}$-blocks are always the same as the $(G \setminus \mathcal{C})$-blocks.

Finally, note that, for $\chi \in \operatorname{Irr}(G)$, $\{\chi\}$ is a $\mathcal{C}$-block if and only if $\chi = \chi^{\mathcal{C}}$ (if $1 \in \mathcal{C}$) or $\chi = \chi^{G \setminus \mathcal{C}}$ (if $1 \notin \mathcal{C}$).

For $b \subseteq \operatorname{Irr}(G)$, we write $(b, \mathcal{C})$ to indicate that b is a $\mathcal{C}$-block. Following [12], we say that there is a *generalized perfect isometry* between two blocks $(b, \mathcal{C})$ and $(b', \mathcal{C}')$ of G and H respectively if there exists a bijection $\mathcal{I}\colon b \to b'$ and signs $\{\eta(\chi),\ \chi \in b\}$ such that, for all χ, $\varphi \in b$, $\langle\, \chi, \varphi \,\rangle_{\mathcal{C}} = \langle\, \eta(\chi)\mathcal{I}(\chi), \eta(\varphi)\mathcal{I}(\varphi) \,\rangle_{\mathcal{C}'}$. Writing $\mathcal{I}_\eta(\chi)$ for $\eta(\chi)\mathcal{I}(\chi)$ ($\chi \in b$), we will say, with a slight abuse of notation, that $\mathcal{I}_\eta\colon (b, \mathcal{C}) \to (b', \mathcal{C}')$ is a generalized perfect isometry. Note that $\mathcal{I}_\eta$ induces an $\mathbb{C}$-vector space isomorphism between the subspaces of $\mathbb{C}\operatorname{Irr}(G)$ and $\mathbb{C}\operatorname{Irr}(H)$ generated by b and b' respectively, which we will also denote by $\mathcal{I}_\eta$.

Our main tool to construct a p-basic set of $\mathfrak{S}_n$ as in Theorem 1.1 will be the following fact (cf Proposition 2.2): with the notations above, if $\mathcal{I}_\eta\colon (b, \mathcal{C}) \to (b', \mathcal{C}')$ is a generalized perfect isometry, then $B \subseteq b$ is a $\mathcal{C}$-basic set of b if and only if $\mathcal{I}(B) \subseteq b'$ is a $\mathcal{C}'$-basic set of b'.

For any odd prime p and any positive integer w, we set

$$G_{p,w} = (\mathbb{Z}_p \rtimes \mathbb{Z}_{p-1}) \wr \mathfrak{S}_w.$$

We can now explain more explicitly our strategy to prove Theorem 1.1.

- We obtain a generalized perfect isometry
$$\mathcal{I}_\eta\colon (\operatorname{Irr}(G_{p,w}), \mathcal{C}_\emptyset) \to (b, p\text{-reg}),$$
where b is any p-block of $\mathfrak{S}_n$ of weight $w > 0$, and $\mathcal{C}_\emptyset$ is a union of conjugacy classes described in §3.2.
- The signature ε of $\mathfrak{S}_n$ permutes the p-blocks of $\mathfrak{S}_n$. We construct a $\mathcal{C}_\emptyset$-basic set $B_\emptyset$ of $G_{p,w}$ such that, if b is an ε-stable p-block of $\mathfrak{S}_n$ of weight $w > 0$, then $\mathcal{I}(B_\emptyset)$ is an ε-stable p-basic set of b.
- We use this to obtain an ε-stable p-basic set of $\mathfrak{S}_n$, and check that Theorem 1.1(2) holds.

The article is organized as follows. In Section 2, we show that a $\mathcal{C}$-basic set of a $\mathcal{C}$-block b is mapped via a generalized perfect isometry $\mathcal{I}_\eta\colon (b, \mathcal{C}) \to (b', \mathcal{C}')$ to a

$\mathcal{C}'$-basic set of b'. In Section 3, we present a generalized perfect isometry, based on the work of the second author in [7], between any p-block of $\mathfrak{S}_n$ of weight $w > 0$ and $\operatorname{Irr}(G_{p,w})$, with respect to a certain union of conjugacy classes $\mathcal{C}_\emptyset$. In Section 4, we describe a $\mathcal{C}_\emptyset$-basic set of $G_{p,w}$. Finally in Section 5, we prove Theorem 1.1. We will show that this p-basic set of $\operatorname{Irr}(\mathfrak{S}_n)$ satisfies Geck's conditions. As a consequence, we prove in Section 5 that, for any odd prime p, there is a p-basic of $\mathfrak{A}_n$. Finally, in Section 6, we give some results on the decomposition numbers of $\mathfrak{A}_n$.

2. RESULTS ON BASIC SETS AND GENERALIZED PERFECT ISOMETRIES

We keep the notation as in Section 1.

Lemma 2.1. *Let G be a finite group and let $\mathcal{C}$ be a union of conjugacy classes of G. We denote by $\operatorname{Bk}_{\mathcal{C}}(G)$ the set of $\mathcal{C}$-blocks of G. If B is a $\mathcal{C}$-basic set of G then, for every $b \in \operatorname{Bk}_{\mathcal{C}}(G)$, $B \cap b$ is a $\mathcal{C}$-basic set of b. Conversely, if every $b \in \operatorname{Bk}_{\mathcal{C}}(G)$ has a $\mathcal{C}$-basic set, B_b say, then*

$$B = \bigcup_{b \in \operatorname{Bk}_{\mathcal{C}}(G)} B_b$$

is a $\mathcal{C}$-basic set of G.

Proof. We suppose that B is a $\mathcal{C}$-basic set of G. Let b be a $\mathcal{C}$-block of G. We will prove that $b \cap B$ is a $\mathcal{C}$-basic set of b. It is clear that $(b \cap B)^{\mathcal{C}}$ is free, being a subset of $B^{\mathcal{C}}$. We have to prove that $(b \cap B)^{\mathcal{C}}$ generates $b^{\mathcal{C}}$ over $\mathbb{Z}$. Let $\chi \in b$. Since B is a $\mathcal{C}$-basic set of G, there are integers $a_{\chi,\psi}$ $(\psi \in B)$ such that

$$\chi^{\mathcal{C}} = \sum_{\psi \in B} a_{\chi,\psi}\, \psi^{\mathcal{C}}.$$

Now take any $\varphi \in \operatorname{Irr}(G)$. If $\varphi \notin b$, then $\langle \chi^{\mathcal{C}}, \varphi \rangle_G = 0$ and $\langle \psi^{\mathcal{C}}, \varphi \rangle_G = 0$ for all $\psi \in B \cap b$, so that $\langle \chi^{\mathcal{C}} - \sum_{\psi \in B \cap b} a_{\chi,\psi}\, \psi^{\mathcal{C}}, \varphi \rangle_G = 0$. On the other hand, if $\varphi \in b$, then $\langle \psi^{\mathcal{C}}, \varphi \rangle_G = 0$ for all $\psi \notin b$, so that $\langle \chi^{\mathcal{C}} - \sum_{\psi \in B \cap b} a_{\chi,\psi}\, \psi^{\mathcal{C}}, \varphi \rangle_G = \langle \chi^{\mathcal{C}} - \sum_{\psi \in B} a_{\chi,\psi}\, \psi^{\mathcal{C}}, \varphi \rangle_G = 0$. Hence we deduce that

$$\chi^{\mathcal{C}} = \sum_{\psi \in B \cap b} a_{\chi,\psi}\, \psi^{\mathcal{C}}.$$

Conversely, it is clear that, if $B = \cup_{b \in \operatorname{Bk}_{\mathcal{C}}(G)} B_b$, then $B^{\mathcal{C}}$ generates $\operatorname{Irr}(G)^{\mathcal{C}}$ over $\mathbb{Z}$. We now show that $B^{\mathcal{C}}$ is free. Suppose there are integers a_ψ $(\psi \in B)$ such that

$$\sum_{\psi \in B} a_\psi\, \psi^{\mathcal{C}} = 0.$$

Let b be a $\mathcal{C}$-block of G. The same argument as previously shows that

$$\sum_{\psi \in B \cap b} a_\psi\, \psi^{\mathcal{C}} = 0.$$

Since $B \cap b = B_b$ and $B_b^{\mathcal{C}}$ is free, we deduce that $a_\psi = 0$ for all $\psi \in b$, and thus for all $\psi \in B$. □

Our main tool is provided by the following

Proposition 2.2. *Let $\mathcal{I}_\eta \colon (b, \mathcal{C}) \to (b', \mathcal{C}')$ be a generalized perfect isometry between two blocks b and b' of finite groups G and H respectively. Then B is a $\mathcal{C}$-basic set of b if and only if the set $B' = \mathcal{I}(B) = \{\eta(\psi)\mathcal{I}_\eta(\psi),\ \psi \in B\}$ is a $\mathcal{C}'$-basic set of b'.*

Proof. Take any $\chi \in b$. We have

$$\chi^{\mathcal{C}} = \sum_{\psi \in \mathrm{Irr}(G)} \langle\, \chi^{\mathcal{C}}, \psi \,\rangle_G \, \psi = \sum_{\psi \in \mathrm{Irr}(G)} \langle\, \chi, \psi \,\rangle_{\mathcal{C}} \, \psi = \sum_{\psi \in b} \langle\, \chi, \psi \,\rangle_{\mathcal{C}} \, \psi = \sum_{\psi \in b} \langle\, \chi^{\mathcal{C}}, \psi \,\rangle_G \, \psi.$$

It follows that

$$\begin{aligned}
\mathcal{I}_\eta(\chi^{\mathcal{C}}) &= \mathcal{I}_\eta\left(\sum_{\psi \in b} \langle\, \chi^{\mathcal{C}}, \psi \,\rangle_G \psi\right) \\
&= \sum_{\psi \in b} \langle\, \chi^{\mathcal{C}}, \psi \,\rangle_G \, \mathcal{I}_\eta(\psi) \\
&= \sum_{\psi \in b} \langle\, \chi, \psi \,\rangle_{\mathcal{C}} \, \mathcal{I}_\eta(\psi) \\
&= \sum_{\psi \in b} \langle\, \mathcal{I}_\eta(\chi), \mathcal{I}_\eta(\psi) \,\rangle_{\mathcal{C}'} \, \mathcal{I}_\eta(\psi) \\
&= \sum_{\psi \in b} \langle\, \mathcal{I}_\eta(\chi), \eta(\psi)\mathcal{I}(\psi) \,\rangle_{\mathcal{C}'} \, \eta(\psi)\, \mathcal{I}(\psi) \\
&= \sum_{\psi' \in b'} \langle\, \mathcal{I}_\eta(\chi), \psi' \,\rangle_{\mathcal{C}'} \, \psi' \\
&= \sum_{\psi' \in b'} \langle\, \mathcal{I}_\eta(\chi)^{\mathcal{C}'}, \psi' \,\rangle_H \, \psi' \\
&= \sum_{\psi' \in \mathrm{Irr}(H)} \langle\, \mathcal{I}_\eta(\chi)^{\mathcal{C}'}, \psi' \,\rangle_H \, \psi' \\
&= \mathcal{I}_\eta(\chi)^{\mathcal{C}'}
\end{aligned}$$

We now suppose that B is a $\mathcal{C}$-basic set of b. Take any $\theta \in b'$. There exists $\chi \in b$ such that $\mathcal{I}_\eta(\chi) = \eta(\chi)\theta$. Since B is a $\mathcal{C}$-basic set of b, there are integers $a_{\chi,\psi}$ ($\psi \in B$) such that

$$\chi^{\mathcal{C}} = \sum_{\psi \in B} a_{\chi,\psi} \psi^{\mathcal{C}}.$$

Using the computation above, we deduce that

$$\eta(\chi)\theta^{\mathcal{C}'} = \mathcal{I}_\eta(\chi^{\mathcal{C}}) = \sum_{\psi \in B} a_{\chi,\psi} \mathcal{I}_\eta(\psi^{\mathcal{C}}) = \sum_{\psi \in B} \underbrace{a_{\chi,\psi}\eta(\psi)}_{\in \mathbb{Z}} \; \underbrace{\mathcal{I}(\psi)^{\mathcal{C}'}}_{\in B'^{\mathcal{C}'}}.$$

Hence the family $B'^{\mathcal{C}'}$ generates $b'^{\mathcal{C}'}$ over $\mathbb{Z}$. We now prove that this family is free. Suppose there are integers b_ψ ($\psi \in B$) such that

$$\sum_{\psi \in B} b_\psi \, \mathcal{I}(\psi)^{\mathcal{C}'} = \sum_{\psi \in B} b_\psi \, \eta(\psi) \, \mathcal{I}_\eta(\psi)^{\mathcal{C}'} = 0.$$

Then we have

$$\mathcal{I}_\eta\left(\sum_{\psi \in B} b_\psi \, \eta(\psi) \, \psi^{\mathcal{C}}\right) = 0.$$

Since $\mathcal{I}_\eta$ is an isomorphism, its kernel is trivial, so that

$$\sum_{\psi \in B} (b_\psi \, \eta(\psi)) \, \psi^{\mathcal{C}} = 0.$$

Now, using the fact that B is a $\mathcal{C}$-basic set of b, we deduce that, for all $\psi \in B$, $b_\psi \eta(\psi) = 0$, and thus $b_\psi = 0$. Hence $B'^{\mathcal{C}'}$ is free, and B' is a $\mathcal{C}'$-basic set of b'.

□

3. GENERALIZED PERFECT ISOMETRY

In this section, we present the generalized perfect isometry we are going to use to reduce our problem from the symmetric group to a wreath product.

3.1. **Some results on partitions.**

Throughout this section, we fix $n \in \mathbb{N}$, p an odd prime, and b a p-block of the symmetric group $\mathfrak{S}_n$. We write $\mathrm{Irr}(\mathfrak{S}_n) = \{\chi_\lambda,\, \lambda \vdash n\}$ as above. By the Nakayama Conjecture (cf [11, 6.1.21]), the irreducible complex characters in b are labelled by the partitions of n with a given p-core, γ say. In particular, we can consider the *weight* w of b, that is the p-weight of any partition of n labelling some irreducible character in b. The characters in b can be parametrized by the set of *p-tuples of partitions* of w, using the abacus. Since we will use this description later on, we present it here on an example. For a complete study, we refer to [11, §2.7] (note however that the abacus we describe here is the horizontal mirror image of that used by James and Kerber).

Consider the lower-right quarter plane (see below). We cut the axes in segments of length 1, and label these by the integers. We choose an arbitrary segment on the vertical axis to be the *origin*, indicated by the symbol $\triangleright$. Whenever we say *symmetric* (respectively reflection), we mean symmetric with respect to (or reflection against) the diagonal (Δ). We label by $\triangledown$ the reflection of the origin (this will play a role in the proof of Lemma 3.1).



Now take for example $p = 3$ and the partition $\lambda = (4, 4, 4, 3, 2)$ of $n = 17$. We put the Young diagram of λ in the upper-left corner of the quarter plane, and consider its (infinite) rim, in bold below.



We see the rim as an infinite sequence of vertical and horizontal dashes of length 1, which, like the axes, we can label by the integers, in such a way that both labellings coincide whenever they meet. In particular, whichever partition we take, exactly one dash in the rim is labelled by the same integer as the origin segment. We say that this dash is *labelled by the origin*, and call it *origin dash*. We can now construct the abacus of λ using this sequence of dashes. We put beads on $p = 3$ runners, going from bottom to top and left to right, putting a bead for each vertical dash and an empty spot for each horizontal one. For this construction to be uniquely defined, we put the bead or empty spot corresponding to the origin dash on the first runner. We get



We see that we thus have a bijection which, to a partition, associates its abacus. If we change the origin, we also change the bijection. But the bijection only depends on the value modulo p of the integer labelling the origin segment. Note that we also get these p different bijections if we fix the origin, but make p different constructions by changing the runner on which we store the origin dash.

The p-information is visible in the abacus in a natural way. For any positive integer m, we call *m-hook* (respectively *(m)-hook*) in a partition any hook of length m (respectively divisible by m). If k is a positive integer, then any kp-rim-hook in λ corresponds to a bead in the abacus which lies, on the same runner, k places above an empty spot. Moreover, this correspondence is bijective.

The removal of a kp-rim-hook in the Young diagram is achieved by moving the corresponding bead down to the empty spot.

By removing all the p-hooks from its Young diagram (which is equivalent to removing all its p-rim-hooks), we get the p-core $\gamma(\lambda)$ of λ, and the corresponding abacus:



To each runner of the abacus, we associate a partition as follows. When we move all the beads as far down as possible, we get one part for each bead that we move, of length the number of places the bead goes down. The resulting p-tuple $\alpha_\lambda = (\lambda^{(1)}, \ldots, \lambda^{(p)})$ of partitions is the *p-quotient* of λ. In the above example, we get $\alpha_\lambda = (\lambda^{(1)}, \lambda^{(2)}, \lambda^{(3)}) = ((1), (2), (1,1))$. The lengths of the $\lambda^{(i)}$'s add up to the p-weight w of λ. We write this as $\alpha_\lambda \Vdash w$. For any positive integer k, we call k-hook in the quotient of λ any k-hook in one of the $\lambda^{(i)}$'s. Then there is a bijection between the set of k-hooks in the p-quotient of λ and the set of kp-hooks in λ.

One sees easily that the partition λ is uniquely determined by its p-core and p-quotient.

We now want to study the effect of conjugation on these descriptions. This is described by the following

Lemma 3.1. *Conjugation of partitions induces a permutation of order 2 on the runners of the p-abacus, which, up to the choice of the origin, is the reflection against the middle runner. With such a choice of origin, if λ has p-quotient $\alpha_\lambda = (\lambda^{(1)}, \ldots, \lambda^{(p)})$, then λ^* has p-quotient $\alpha_{\lambda^*} = (\lambda^{(p)^*}, \ldots, \lambda^{(1)^*})$.*

Proof. Take any partition λ. Throughout the proof, we use λ as above as an example. The rim of λ^* is visible in the Young diagram of λ: we just need to exchange the roles of horizontal and vertical dashes in the rim of λ. In terms of abacus, this means replacing each bead in the abacus of λ by an empty spot, and each empty spot by a bead. This corresponds to transforming the partition stored on each runner into its conjugate. If we then turn this new abacus upside down (that is, rotate it 180 degrees), we obtain **an** abacus which represents λ^*. In our example, we get



which then becomes



However, this is **the** abacus of λ^* (in the bijection we mentionned before) only if the position labelled by the origin $\triangleright$ is on the first runner. The spot that was labelled by $\triangleright$ in the abacus of λ is now on the last runner, and labelled by the symmetrized $\triangledown$ of $\triangleright$ on the horizontal axis. Hence the spot now labelled by $\triangleright$ is on the first runner if and only if the number of spots which separate it from the spot labelled by $\triangledown$ is $p-1$ plus a multiple of p. But this number is exactly the distance, following the axes, between $\triangledown$ and $\triangleright$. Hence our construction gives **the** abacus of λ^* if and only if this distance is congruent to -1 modulo p (as is the case in our example, since $11 = -1 \ (mod\ 3)$). If this is the case, then, from the description we gave above, we see immediately that, if λ has p-quotient $\alpha_\lambda = (\lambda^{(1)}, \ldots, \lambda^{(p)})$, then λ^* has p-quotient $\alpha_{\lambda^*} = (\lambda^{(p)^*}, \ldots, \lambda^{(1)^*})$ (in our example, we get that $\alpha_{\lambda^*} = (\lambda^{(3)^*}, \lambda^{(2)^*}, \lambda^{(1)^*}) = ((2), (1,1), (1)))$. $\square$

This leads to the following

Convention 3.2. *We choose the origin so that the permutation of Lemma 3.1 is the reflection against the* r*-th runner, where* $r = (p+1)/2$.

Remark 3.3. *Note that, with Convention 3.2, applying the transformation described in the proof of Lemma 3.1 to the abacus of* λ *always gives the abacus of* λ^*. *However, it corresponds to* $(\gamma(\lambda), \alpha_\lambda) \longmapsto (\gamma(\lambda)^*, \alpha_{\lambda^*})$, *and it's easy to see that* $\gamma(\lambda)^* = \gamma(\lambda^*)$. *In particular, this transformation needs not preserve the core* $\gamma(\lambda)$, *and thus the block in which lies the character* χ_λ. *If* $\gamma(\lambda) \neq \gamma(\lambda^*)$, *then there still is a character* χ_μ *in the same* p*-block as* χ_λ *and such that* $\alpha_\mu = \alpha_{\lambda^*}$, *but* $\mu \neq \lambda^*$.

We also include here a lemma about the partition $\overline{\lambda}$ defined in Section 1 for a self-conjugate partition, and which will be useful later.

Lemma 3.4. *Let* p *be an odd prime,* $r = (p+1)/2$, *and let* λ *be a self-conjugate partition of* n. *Then the partition* $\overline{\lambda}$ *is* p*-regular, if and only if the* r*-th part of the quotient* α_λ *of* λ *is empty.*

Proof. A hook in the Young diagram of a partition is called *diagonal* if its top left box lies on the diagonal (Δ). For any self-conjugate $\lambda \vdash n$, the parts of $\overline{\lambda}$ are therefore exactly the lengths of the diagonal hooks in λ.

Now, by conjugation, any (p)-hook stored on the i-th runner of the abacus of (any) λ is tranformed into its reflection, and is stored on the $(p+1-i)$-th runner in the abacus of λ^*. Thus the hooks on the i-th runner of λ are exactly the reflections of those on the $(p+1-i)$-th runner of λ^*.

In particular, if λ is self-conjugate, then $\alpha_\lambda = \alpha_{\lambda^*}$, and each of the hooks on the r-th runner is its own reflection (and all the others move). Hence the r-th runner in the abacus of λ stores exactly the symmetric (p)-hooks, which are precisely the diagonal (p)-hooks. Since these correspond to the parts divisible by p in $\overline{\lambda}$, we easily get the result.

Remark: more generally, even if λ is not self-conjugate, one can prove that the r-th runner in the abacus of λ stores exactly the diagonal (p)-hooks of λ.

□

Remark 3.5. *In particular, if* $w = 0$, *then* λ *has no* (p)*-hook (and, a fortiori, no diagonal* (p)*-hook), so that* $\overline{\lambda}$ *is always* p*-regular.*

3.2. **Wreath product.** We denote by $\mathbb{Z}_p$ and $\mathbb{Z}_{p-1}$ the cyclic groups of order p and $p-1$ respectively. Note that $\mathbb{Z}_{p-1} \cong \operatorname{Aut}(\mathbb{Z}_p)$, so that we can construct the semidirect product $N = \mathbb{Z}_p \rtimes \mathbb{Z}_{p-1}$. Considering $\mathbb{Z}_p$ as the subgroup of $\mathfrak{S}_p$ generated by a p-cycle, we then have $N = N_{\mathfrak{S}_p}(\mathbb{Z}_p)$. We denote by $G_{p,w}$ the wreath product $N \wr \mathfrak{S}_w$. That is, $G_{p,w}$ is the semidirect product $N^w \rtimes \mathfrak{S}_w$, where $\mathfrak{S}_w$ acts by permutation on the w copies of N. Note that, if $w < p$, then $P \cong \mathbb{Z}_p^w$ is a Sylow p-subgroup of $\mathfrak{S}_{pw}$, and $G_{p,w} \cong N_{\mathfrak{S}_{pw}}(P)$.

For a complete description of wreath products and their representations, we refer to [11, Chapter 4].

We write $r = (p+1)/2$. We have $\operatorname{Irr}(N) = \{\psi_1, \ldots, \psi_p\}$, with $\psi_i(1) = 1$ for $1 \leq i \leq p$ and $i \neq r$, and $\psi_r(1) = p-1$. More precisely, writing $1_{\mathbb{Z}_p}$ for the trivial character of $\mathbb{Z}_p$, we have

$$\mathrm{Ind}_{\mathbb{Z}_p}^{N}(1_{\mathbb{Z}_p}) = \sum_{\substack{i=1\\ i\neq r}}^{p} \psi_i \ \text{ and } \ \mathrm{Res}_{\mathbb{Z}_p}^{N}(\psi_i) = 1_{\mathbb{Z}_p} \ \ (1 \leq i \leq p,\, i \neq r),$$

and

$$\mathrm{Res}_{\mathbb{Z}_p}^{N}(\psi_r) = \eta_2 + \cdots + \eta_p \ \text{ and } \ \mathrm{Ind}_{\mathbb{Z}_p}^{N}(\eta_i) = \psi_r \ \ (2 \leq i \leq p),$$

where $\{\eta_2, \ldots, \eta_p\} = \mathrm{Irr}(\mathbb{Z}_p) \setminus \{1_{\mathbb{Z}_p}\}$.

The irreducible complex characters of $G_{p,w} = N \wr \mathfrak{S}_w$ are parametrized by the p-tuples of partitions of w as follows. Take any $\alpha = (\alpha^1, \ldots, \alpha^p) \Vdash w$ and consider the irreducible character $\prod_{i=1}^{p} \psi_i^{|\alpha^i|}$ of the *base group* N^w. It can be extended in a natural way to its inertia subgroup $N \wr \mathfrak{S}_{|\alpha^1|} \times \cdots \times N \wr \mathfrak{S}_{|\alpha^p|}$, giving the irreducible character $\prod_{i=1}^{p} \widetilde{\psi_i^{|\alpha^i|}}$ [11, p. 154]. Any extension is of the form $\prod_{i=1}^{p}(\widetilde{\psi_i^{|\alpha^i|}} \otimes \varphi_{\alpha_i})$, with $\varphi_{\alpha_i} \in \mathrm{Irr}(\mathfrak{S}_{|\alpha^i|})$ $(1 \leq i \leq p)$. Then $\chi^\alpha := \mathrm{Ind}_{\prod_{i=1}^{p} N \wr \mathfrak{S}_{|\alpha^i|}}^{N \wr \mathfrak{S}_w}(\prod_{i=1}^{p} \widetilde{\psi_i^{|\alpha^i|}} \otimes \varphi_{\alpha^i}) \in \mathrm{Irr}(G_{p,w})$. Different $\alpha \Vdash w$ give different irreducible characters of $G_{p,w}$, and any irreducible character of $G_{p,w}$ can be obtained in this way.

Note for future reference that, in the above notation, if $\alpha^r = \emptyset$, then $P = \mathbb{Z}_p^w \subseteq \ker(\chi^\alpha)$. Indeed, for all $1 \leq i \leq p$, $i \neq r$, we have $\mathrm{Res}_{\mathbb{Z}_p}^{N}(\psi_i) = 1_{\mathbb{Z}_p}$; thus $\mathrm{Res}_{\mathbb{Z}_p^{|\alpha^i|}}^{N^{|\alpha^i|}}(\psi_i^{|\alpha^i|}) = 1_{\mathbb{Z}_p^{|\alpha^i|}}$, so that $\widetilde{\psi_i^{|\alpha^i|}}(g) = \widetilde{\psi_i^{|\alpha^i|}}(1)$ for all $g \in \mathbb{Z}_p^{|\alpha^i|}$, and $(\widetilde{\psi_i^{|\alpha^i|}} \otimes \varphi_{\alpha_i})(g) = (\widetilde{\psi_i^{|\alpha^i|}} \otimes \varphi_{\alpha_i})(1)$ for all $g \in \mathbb{Z}_p^{|\alpha^i|}$. Since $P \leq N^w \lhd N \wr \mathfrak{S}_w$, we easily get that, for all $g \in P = \mathbb{Z}_p^w$, $\chi^\alpha(g) = \chi^\alpha(1)$.

The conjugacy classes of $G_{p,w}$ are also parametrized by the p-tuples of partitions of w. Let $g_1, \ldots, g_p$ be representatives for the conjugacy classes of N. Note that, when seen as a subgroup of $\mathfrak{S}_p$, N has a unique conjugacy class of p-cycles, for which we take representative g_p. The elements of $G_{p,w} = N \wr \mathfrak{S}_w$ are of the form $(h, \gamma) = ((h_1, \ldots, h_w), \sigma)$, with $h_1, \ldots, h_w \in N$ and $\sigma \in \mathfrak{S}_w$. For any such element, and for any k-cycle $\kappa = (j, j\kappa, \ldots, j\kappa^{k-1})$ in σ, we define the *cycle product* of (h, σ) and κ by

$$g((h, \sigma);\, \kappa) = h_j h_{j\kappa^{-1}} h_{j\kappa^{-2}} \ldots h_{j\kappa^{-(k-1)}}.$$

In particular, $g((h, \sigma);\, \kappa) \in N$. If σ has cycle structure π say, then we form p partitions $(\pi_1, \ldots, \pi_p)$ from π as follows: any cycle κ in π gives a cycle of the same length in π_i if the cycle product $g((h, \sigma);\, \kappa)$ is conjugate to g_i in N. The resulting p-tuple of partitions of w describes the *cycle structure* of (h, γ). Two elements of $G_{p,w}$ are conjugate if and only if they have the same cycle structure.

Define $\mathcal{C}_\emptyset$ to be the set of elements of $G_{p,w}$ with cycle structure $(\pi_1, \ldots, \pi_p)$ such that $\pi_p = \emptyset$.

Note that, in the natural embedding of $G_{p,w}$ in $\mathfrak{S}_{pw}$, for any k-cycle κ of σ, each m-cycle of the cycle product $g((h, \sigma);\, \kappa) \in \mathfrak{S}_p$ corresponds to $gcd(m, k)$ cycles of length $lcm(m, k)$ in $(h, \sigma)\mathfrak{S}_{pw}$. In particular, if $w < p$, then $\mathcal{C}_\emptyset$ is the set of p-regular elements of $G_{p,w}$.

3.3. **A generalized perfect isometry.** We now come to the generalized perfect isometry presented by the second auther in [7]. We first introduce one last piece of notation ([7, Prop. 3.8]). Let $\alpha = (\alpha^1, \ldots, \alpha^p)$ be any p-tuple of partitions of w, and $r = (p+1)/2$ as before. We define $\tilde{\alpha} \Vdash w$ by

$$\tilde{\alpha} = (\alpha^1, \ldots, \alpha^{r-1}, (\alpha^r)^*, \alpha^{r+1}, \ldots, \alpha^p),$$

where * denotes conjugation of partitions.

Note that $\tilde{}$ is a bijection from $\{(\alpha^1, \ldots, \alpha^p) \Vdash w\}$ onto itself, and is the identity on $\{(\alpha^1, \ldots, \alpha^p) \Vdash w, \, \alpha^r = \emptyset\}$.

The main result in [7] can, in our context, be stated as follows

Theorem 3.6. *([7], Theorem 4.1) Let $n \geq 1$ be an integer and p be a prime, and let b be a p-block of weight $w \neq 0$. For any $\chi_\lambda \in b$, denote by α_λ the p-quotient of λ. Then, with the above notations, the bijection*

$$\begin{array}{rcl} \mathcal{J}\colon b & \longrightarrow & \mathrm{Irr}(G_{p,w}) \\ \chi_\lambda & \longmapsto & \chi^{\widetilde{\alpha_\lambda}} \end{array}$$

induces a generalized perfect isometry $\mathcal{J}_\eta\colon (b, p\text{-}reg) \to (\mathrm{Irr}(G_{p,w}), \mathcal{C}_\emptyset)$.

Proof. We give here the main lines of the proof, first for the comfort of the reader, but also because, in fact, Theorem 4.1 in [7] is not stated as above; it is only stated in the case where b is the principal p-block of $\mathfrak{S}_n$, and when $w < p$. However, for our purpose, and for the version of this theorem we give, these two hypotheses are not necessary, as we will try and explain below. The object of [7] is to give an analogue, for the ℓ-blocks of $\mathfrak{S}_n$ (cf [12]), where ℓ is an arbitrary integer, of Broué's Abelian Defect Conjecture (hence the hypotheses on b and w). In particular, in this case, and when $\ell = p$, $\mathcal{C}_\emptyset$ is the set of p-regular elements of $G_{p,w}$. This fails when $w \geq p$, but the generalized perfect isometry we mentionned above still exists.

We consider a p-cycle ω in $\mathfrak{S}_p$ and $L = \langle \omega \rangle$, and we let $N = N_{\mathfrak{S}_p}(L)$. Then $N = L \rtimes \mathrm{Aut}(L) \cong \mathbb{Z}_p \rtimes \mathbb{Z}_{p-1}$, and direct computation shows that N has a single p-block.

Now take any p-block b of weight $w \neq 0$ of $\mathfrak{S}_n$. Take representatives $(h_i = \omega^{i-1}, \, 1 \leq i \leq p)$ for the conjugacy classes of L. As above, we have that the conjugacy classes of $L \wr \mathfrak{S}_w$ are parametrized by the p-tuples $(\pi_1, \ldots, \pi_p) \Vdash w$ of partitions of w. For π_1 in $\mathcal{P}_{\leq w}$ (the set of partitions of at most w), we write $\mathcal{D}_{\pi_1}$ for the set of elements whose cycle structure has first part π_1. In particular, $\mathcal{D}_\emptyset$ is the set of *regular* elements described by Osima (cf [13]). We then have

$$L \wr \mathfrak{S}_w = \coprod_{\pi_1 \in \mathcal{P}_{\leq w}} \mathcal{D}_{\pi_1}.$$

Write $\mathrm{Irr}(L \wr \mathfrak{S}_w) = \{\xi^\alpha, \, \alpha \Vdash w\}$. One fundamental result we use (cf [13] and [12]), and which does not depend on $w \neq 0$, is that, with the notations above, $\lambda \longmapsto \alpha_\lambda$ induces a generalized perfect isometry $\mathcal{I}_\epsilon\colon (b, p\text{-reg}) \to (\mathrm{Irr}(L \wr \mathfrak{S}_w), \mathcal{D}_\emptyset)$. We therefore have, for all $\chi_\lambda, \chi_\mu \in b$,

$$\langle \chi_\lambda, \chi_\mu \rangle_{p\text{-reg}} = \langle \epsilon(\lambda)\xi^{\alpha_\lambda}, \epsilon(\mu)\xi^{\alpha_\mu} \rangle_{\mathcal{D}_\emptyset}. \tag{1}$$

The biggest part of the proof now consists in transporting this to the group $G_{p,w}$ the elements of $\mathcal{C}_\emptyset$. The idea is to work with *singular* elements instead of regular elements, and use induction on w. For any $\alpha, \beta \Vdash w$, we have

$$\langle \xi^\alpha, \xi^\beta \rangle_{\mathcal{D}_\emptyset} = \delta_{\alpha\beta} - \langle \xi^\alpha, \xi^\beta \rangle_{L \wr \mathfrak{S}_w \setminus \mathcal{D}_\emptyset}$$

and

$$\langle \xi^\alpha,\, \xi^\beta \rangle_{L \wr \mathfrak{S}_w \setminus \mathcal{D}_\emptyset} = \sum_{\emptyset \neq \pi_1 \in \mathcal{P}_{\leq w}} \langle \xi^\alpha,\, \xi^\beta \rangle_{\mathcal{D}_{\pi_1}}.$$

The induction on w will be based on an analogue of the Murnaghan-Nakayama Rule which holds in wreath products (cf [14, 4.4]): writing, for any $\emptyset \neq \pi_1 \in \mathcal{P}_{\leq w}$, $\mathcal{D}_{(\pi_1, \ldots, \pi_p)}$ for the conjugacy class of cycle structure $(\pi_1, \ldots, \pi_\ell)$ of $L \wr \mathfrak{S}_w$, and $\mathcal{D}^{w-|\pi_1|}_{(\pi_2, \ldots, \pi_p)}$ for the set of elements of cycle structure $(\emptyset, \pi_2, \ldots, \pi_p)$ in $L \wr \mathfrak{S}_{w-|\pi_1|}$, we have

$$\xi^\alpha(\mathcal{D}_{(\pi_1, \ldots, \pi_p)}) = \sum_{1 \leq s_1, \ldots, s_i \leq p} \ \sum_{\hat{\alpha} \in \mathcal{L}^{\pi_1}_{\alpha, (s_1, \ldots, s_i)}} (-1)^{L_{\alpha\hat{\alpha}}} \xi^{\hat{\alpha}}(\mathcal{D}^{w-|\pi_1|}_{(\pi_2, \ldots, \pi_p)}),$$

where $\mathcal{L}^{\pi_1}_{\alpha, (s_1, \ldots, s_i)}$ is the set of p-tuples of partitions of $w - |\pi_1|$ which can be obtained from α by removing successively a k_1-hook from α^{s_1}, then a k_2-hook from the "s_2-th coordinate" of the resulting p-tuple of partitions of $w - k_1$, etc, and finally a k_i-hook from the "s_i-th coordinate" of the resulting p-tuple of partitions of $w - (k_1 + \cdots + k_{i-1})$, and, for $\hat{\alpha} \in \mathcal{L}^{\pi_1}_{\alpha, (s_1, \ldots, s_i)}$, $L_{\alpha\hat{\alpha}}$ is the sum of the leg-lengths of the hooks removed to get from α to $\hat{\alpha}$.

Writing b_k for the number of k-cycles in π_1, and $d_{\pi_1} = \prod_{1 \leq k \leq w} b_k!(kp)^{b_k}$, we then get, writing $\mathbf{s} = (s_1,\, \ldots,\, s_i)$,

$$\langle \xi^\alpha,\, \xi^\beta \rangle_{\mathcal{D}_{\pi_1}} = \frac{1}{d_{\pi_1}} \sum_{\substack{1 \leq \mathbf{s} \leq p \\ 1 \leq \mathbf{t} \leq p}} \ \sum_{\substack{\hat{\alpha} \in \mathcal{L}^{\pi_1}_{\alpha, \mathbf{s}} \\ \hat{\beta} \in \mathcal{L}^{\pi_1}_{\beta, \mathbf{t}}}} (-1)^{L_{\alpha\hat{\alpha}}} (-1)^{L_{\beta\hat{\beta}}} \langle \xi^{\hat{\alpha}},\, \xi^{\hat{\beta}} \rangle_{\mathcal{D}^{w-|\pi_1|}_\emptyset}.$$

We will use this formula to exhibit a generalized perfect isometry between $\mathrm{Irr}(L \wr \mathfrak{S}_w)$ and $\mathrm{Irr}(N \wr \mathfrak{S}_w)$.

Take $(g_1 = 1,\, \ldots,\, g_p = \omega)$ representatives for the conjugacy classes of N. We have

$$N \wr \mathfrak{S}_w = \coprod_{\pi_p \in \mathcal{P}_{\leq w}} \mathcal{C}_{\pi_p},$$

where $\mathcal{C}_{\pi_p}$ denotes the set of elements whose cycle structure has p-th part π_p, and, for any $\alpha,\, \beta \Vdash w$,

$$\langle \chi^\alpha,\, \chi^\beta \rangle_{N \wr \mathfrak{S}_w \setminus \mathcal{C}_\emptyset} = \sum_{\emptyset \neq \pi_p \in \mathcal{P}_{\leq w}} \langle \chi^\alpha,\, \chi^\beta \rangle_{\mathcal{C}_{\pi_p}}.$$

Now fix any $\emptyset \neq \pi_p \in \mathcal{P}_{\leq w}$, and write b'_k for the number of k-cycles in π_p, and $c_{\pi_p} = \prod_{1 \leq k \leq w} b'_k!(kp)^{b'_k}$. In particular, if $\pi_1 = \pi_p$, then $d_{\pi_1} = c_{\pi_p}$. We then have, by repeated use of the Murnaghan-Nakayama Rule, and with similar notations as above,

$$\langle \chi^\alpha,\, \chi^\beta \rangle_{\mathcal{C}_{\pi_p}} = \frac{1}{c_{\pi_p}} \sum_{\substack{1 \leq \mathbf{s} \leq p \\ 1 \leq \mathbf{t} \leq p}} \ \sum_{\substack{\hat{\alpha} \in \mathcal{L}^{\pi_p}_{\alpha, \mathbf{s}} \\ \hat{\beta} \in \mathcal{L}^{\pi_p}_{\beta, \mathbf{s}}}} \Psi_{\mathbf{s}}(g_p) \overline{\Psi_{\mathbf{t}}(g_p)} (-1)^{L_{\alpha\hat{\alpha}}} (-1)^{L_{\beta\hat{\beta}}} \langle \chi^{\hat{\alpha}},\, \chi^{\hat{\beta}} \rangle_{\mathcal{C}^{w-|\pi_p|}_\emptyset},$$

where $\Psi_{\mathbf{s}}(g_p) = \psi_{s_1}(g_p) \ldots \psi_{s_i}(g_p)$ (with $\mathrm{Irr}(N) = \{\psi_1,\, \ldots,\, \psi_p\}$). Now the transformation $\hat{}$ defined before Theorem 3.6 is introduced precisely to get rid of the factor $\Psi_{\mathbf{s}}(g_p)$ (which is only problematic when $\mathbf{s}$ contains $r = (p+1)/2$, since

$\psi_r(\omega) = -1$ but $\psi_i(\omega) = 1$ for $i \neq r$). For $\alpha \Vdash w$, we let $\chi_0^\alpha = (-1)^{|\alpha_r|}\chi^{\hat{\alpha}}$, and we can use induction on w to prove that, for any α, $\beta \Vdash w$,

$$\langle \chi_0^\alpha, \chi_0^\beta \rangle_{\mathcal{C}_\emptyset} = \langle \xi^\alpha, \xi^\beta \rangle_{\mathcal{D}_\emptyset} \tag{2}$$

(cf [7, 3.10]). Note that the result is clearly true when $w = 0$ since $\mathcal{C}_\emptyset = \mathcal{D}_\emptyset = \emptyset$ and $\chi^\emptyset = \xi^\emptyset = 1$. Composing the results of (1) and (2), we get the desired generalized perfect isometry. $\square$

Note that the above result does not depend on the order in which we list the irreducible characters of N (provided we modify the definition of $\tilde{}$ in accordance), or on the runner of the abacus we choose to be fixed by conjugation. Each choice of labelling for the characters of N and for the fixed runner will give an (a priori different) generalized perfect isometry. In particular, it was legitimate to make the choices we made.

4. A $\mathcal{C}_\emptyset$-BASIC SET FOR $G_{p,w}$

We now want to exhibit a $\mathcal{C}_\emptyset$-basic set for $G_{p,w}$, which we will then transport to $\mathfrak{S}_n$ using the generalized perfect isometry given by Theorem 3.6. Our main tool to do so will be Proposition 4.2. In the course of its proof, we will need the following general lemma about semidirect products, which we therefore state separately.

Lemma 4.1. *Let $G = H \rtimes K$ be a finite group, and $\pi\colon G \longrightarrow K$ be the canonical surjection. If $k \in K$ is conjugate in G to $g \in G$, then k is conjugate in K to $\pi(g)$. In particular, two elements of K are conjugate in G if and only if they are conjugate in K.*

Proof. Suppose $g \in G$ and $k \in K$ are conjugate in G. Then there exist $h \in H$ and $\ell \in K$ such that $g = (h\ell)k(h\ell)^{-1}$. Thus $g = h(\ell k \ell^{-1})h^{-1}$, and $\ell k \ell^{-1} = k' \in K$. Hence $g = (h(h^{-1})^{k'})k'$, and, by uniqueness of the (H, K)-decomposition in G, we must have $k' = \pi(g)$, so that k and $\pi(g)$ are conjugate in K. $\square$

Proposition 4.2. *Suppose $G = H \rtimes K$ is a finite group, and $\pi\colon G \longrightarrow K$ is the canonical surjection. Suppose $\mathcal{C}$ is a union of conjugacy classes of G such that $K \subseteq \mathcal{C}$ and $|Cl_G(\mathcal{C})| = |Cl_K(K)|$ (i.e. the numbers of G-conjugacy classes in $\mathcal{C}$ and of conjugacy classes in K are the same). Then $\{\psi \circ \pi,\ \psi \in \mathrm{Irr}(K)\}$, the set of irreducible characters of G with H in their kernel, is a $\mathcal{C}$-basic set for G.*

Note that Proposition 4.2 is known in the case where H is a p-group, K is a p'-group, and $\mathcal{C}$ is the set of p-regular elements of G.

Proof. Let $\mathcal{B} = \{\psi \circ \pi,\ \psi \in \mathrm{Irr}(K)\}$. Since $|Cl_G(\mathcal{C})| = |Cl_K(K)| = |\mathcal{B}|$, it suffices to prove that $\mathcal{B}^{\mathcal{C}}$ generates $\mathrm{Irr}(G)^{\mathcal{C}}$ over $\mathbb{Z}$.

Since $\mathcal{K} \subseteq \mathcal{C}$, and since, by Lemma 4.1, any two elements of K are conjugate in K if and only if they are conjugate in G, we get that $|Cl_K(K)| = |\{Cl_G(g) \in Cl_G(\mathcal{C})$ such that $\exists g_K \in K,\ Cl_G(g) = Cl_G(g_K)\}|$. Since $|Cl_G(\mathcal{C})| = |Cl_K(K)|$, this shows that every G-conjugacy class in $\mathcal{C}$ has a representative in K.

Now take any $\chi \in \mathrm{Irr}(G)$, and write $\mathrm{Res}_K^G(\chi) = \sum_{\psi \in \mathrm{Irr}(K)} a_{\chi\psi}\psi$, with $a_{\chi\psi} \in \mathbb{N}$ for all $\psi \in \mathrm{Irr}(K)$. Take any $g \in \mathcal{C}$. By the above, there exists $g_K \in K$ such that $Cl_G(g) = Cl_G(g_K)$, and, by Lemma 4.1, g_K and $\pi(g)$ are conjugate in K. We therefore get

$$\chi(g) = \chi(g_K) = \sum_{\psi \in \mathrm{Irr}(K)} a_{\chi\psi}\psi(g_K) = \sum_{\psi \in \mathrm{Irr}(K)} a_{\chi\psi}\psi(\pi(g))$$

(the first equality being true because g and g_K are conjugate in G, the last because g_K and $\pi(g)$ are conjugate in K). Since this holds for any $g \in \mathcal{C}$, we finally get that $\chi^{\mathcal{C}} = \sum_{\psi \in \mathrm{Irr}(K)} a_{\chi\psi}(\psi \circ \pi)^{\mathcal{C}}$, so that $\mathcal{B}^{\mathcal{C}}$ generates $\mathrm{Irr}(G)^{\mathcal{C}}$ over $\mathbb{Z}$.

□

We can now prove the main result of this section

Theorem 4.3. *The set* $B_{\emptyset} = \{\chi^{\alpha} \in \mathrm{Irr}(G_{p,w}) \mid \alpha = (\alpha^1, \ldots, \alpha^p) \Vdash w, \alpha^r = \emptyset\}$ *is a* $\mathcal{C}_{\emptyset}$*-basic set for* $G_{p,w}$*.*

Proof. We start by proving that we can somehow apply Proposition 4.2 to $G_{p,w}$ to obtain a $\mathcal{C}_{\emptyset}$-basic set. Write $G_{p,w} = (\mathbb{Z}_p \rtimes \mathbb{Z}_{p-1}) \wr \mathfrak{S}_w$ as

$$\{(((a_1, \ldots, a_w), (b_1, \ldots, b_w)), \sigma),\ a_i \in \mathbb{Z}_p,\ b_i \in \mathbb{Z}_{p-1}\ (1 \leq i \leq w),\ \sigma \in \mathfrak{S}_w\}.$$

One proves easily, using the properties of semidirect products and the fact that $\mathfrak{S}_w$ acts by permutation on the w copies of $\mathbb{Z}_p \rtimes \mathbb{Z}_{p-1}$, the following:

- $\mathcal{H} = \{(((a_1, \ldots, a_w), (1, \ldots, 1)), 1)\}$ is a normal subgroup of $G_{p,w}$, isomorphic to $\mathbb{Z}_p^w$,
- $\{(((1, \ldots, 1), (1, \ldots, 1)), \sigma)\}$ is a subgroup of $G_{p,w}$, isomorphic to $\mathfrak{S}_w$,
- $\mathcal{K} = \{(((1, \ldots, 1), (b_1, \ldots, b_w)), \sigma)\}$ is a subgroup of $G_{p,w}$, isomorphic to $\mathbb{Z}_{p-1} \wr \mathfrak{S}_w$,
- $\mathcal{H} \cap \mathcal{K} = \{1\}$, and $\mathcal{H}\mathcal{K} = G_{p,w}$.

We can therefore write $G_{p,w}$ as a semidirect product $\mathcal{H} \rtimes \mathcal{K}$.

For any $k = (((1, \ldots, 1), (b_1, \ldots, b_w)), \sigma) \in \mathcal{K}$, we see that all cycle products of k are products of some b_i's, hence belong to $\mathbb{Z}_{p-1}$, and, in particular, are not conjugate to the p-cycle of $N = \mathbb{Z}_p \rtimes \mathbb{Z}_{p-1}$. Thus $\mathcal{K} \subseteq \mathcal{C}_{\emptyset}$.

Finally, the numbers of conjugacy classes in $\mathcal{C}_{\emptyset}$ and in $\mathcal{K}$ are the same, since both of these sets can be labelled by the $(p-1)$-tuples of partitions of w. We therefore deduce from Proposition 4.2 that the irreducible characters of $G_{p,w}$ with $\mathcal{H}$ in their kernel form a $\mathcal{C}_{\emptyset}$-basic set for $G_{p,w}$.

Now, as we remarked when we presented the irreducible characters of $G_{p,w}$, if $\alpha = (\alpha^1, \ldots, \alpha^p) \Vdash w$ is such that $\alpha^r = \emptyset$, then $\chi^{\alpha} \in \mathrm{Irr}(G_{p,w})$ has $\mathcal{H}$ in its kernel. Since these characters can naturally be labelled by the $(p-1)$-tuples of partitions of w, they must be all the irreducible characters of $G_{p,w}$ with $\mathcal{H}$ in their kernel. This ends the proof.

□

5. A p-BASIC SET FOR $\mathfrak{A}_n$

5.1. **Proof of Theorem 1.1.** We keep the notation as above. As previously, for any positive integer w and odd prime p, we set $G_{p,w} = (\mathbb{Z}_p \rtimes \mathbb{Z}_{p-1}) \wr \mathfrak{S}_w$. Using Lemma 2.1 we first reduce the problem as follows. The signature ε induces a permutation of order 2 on the set $\mathrm{Bk}_p(\mathfrak{S}_n)$ of p-blocks of $\mathfrak{S}_n$. Each orbit has one or two p-blocks (which then have the same weight). Suppose that the orbit $\omega = \{b, \varepsilon b\}$ has two p-blocks of weight $w \geq 0$. If $w = 0$, then b and εb each consist of a single character (which vanishes on p-singular elements), so that $b \cup \varepsilon b$ is a p-basic set for itself, which satisfies Condition (2) of Theorem 1.1 by Remark 3.5. Suppose then that $w \neq 0$. By Theorem 3.6 there is a generalized perfect isometry

$$\mathcal{I}_\eta \colon (\mathrm{Irr}(G_{p,w}), \mathcal{C}_\emptyset) \to (b, p\text{-reg})$$

($\mathcal{I}_\eta$ is the inverse of the isometry $\mathcal{J}_\eta$ given by Theorem 3.6). Furthermore, by Theorem 4.3, we know that $G_{p,w}$ has a $\mathcal{C}_\emptyset$-basic set $B_\emptyset$. Hence we deduce, using Proposition 2.2, that b has a p-basic set $B'_\emptyset$. Consider now the set $\varepsilon B'_\emptyset \subseteq \varepsilon b$. This is clearly a p-basic set of εb. Moreover, by Lemma 2.1, the set $B'_\emptyset \cup \varepsilon B'_\emptyset$ is a p-basic set of $b \cup \varepsilon b$, and it is ε-stable by construction. We now remark that, if $\chi_\lambda \in b \cup \varepsilon b$, then λ is never self-conjugate, because the p-core of a self-conjugate partition is also self-conjugate. Hence Condition (2) of Theorem 1.1 is automatically satisfied by $B'_\emptyset \cup \varepsilon B'_\emptyset$.

We can proceed as above for every orbit of size two, and we therefore obtain a p-basic set of

$$\bigcup_{\omega = \{b, \varepsilon b\},\, |\omega| = 2} b \cup \varepsilon b,$$

which satisfies the conditions of Theorem 1.1. Hence to prove Theorem 1.1, it is sufficient to prove that every ε-stable p-block of $\mathfrak{S}_n$ has a p-basic set satisfying the conditions of Theorem 1.1.

Let b be an ε-stable p-block of $\mathfrak{S}_n$ of weight $w \geq 0$. As above, if $w = 0$, then b is a p-basic set for itself, and satisfies Condition (2) of Theorem 1.1 by Remark 3.5. We therefore suppose that $w \neq 0$. As in the previous case, write $\mathcal{I}_\eta \colon (\mathrm{Irr}(G_{p,w}), \mathcal{C}_\emptyset) \to (b, p\text{-reg})$ for the (inverse of the) generalized perfect isometry given by Theorem 3.6. Let be the $\mathcal{C}_\emptyset$-basic set of $G_{p,w}$ obtained in Theorem 4.3. By Proposition 2.2, $B'_\emptyset = \mathcal{I}(B_\emptyset)$ is a p-basic set of b. It follows from the definition of $\mathcal{I}$ that

$$B'_\emptyset = \{\chi_\lambda \in b \mid \alpha_\lambda = (\alpha^1, \ldots, \alpha^p) \Vdash w,\ \alpha^{(p+1)/2} = \emptyset\}.$$

Let $\chi_\lambda \in B'_\emptyset$. Then, by Lemma 3.1 and Convention 3.2, the r-th part ($r = (p+1)/2$) of the p-quotient of λ^* is empty. Hence $\varepsilon\chi_\lambda = \chi_{\lambda^*} \in B'_\emptyset$. This proves that $B'_\emptyset$ is ε-stable. Lemma 3.4 immediately implies that $B'_\emptyset$ satisfies Condition (2) of Theorem 1.1. This proves the result.

5.2. **A p-basic set for $\mathfrak{A}_n$.** We first recall the parametrizations of the classes and irreducible characters of $\mathfrak{A}_n$. Because of Clifford's theory [2, III.2.12], we know that the set of irreducible characters of $\mathfrak{A}_n$ consists of the irreducible constituents occurring in the restrictions to $\mathfrak{A}_n$ of irreducible characters of $\mathfrak{S}_n$. More precisely, let χ_λ be an irreducible character of $\mathfrak{S}_n$. If $\lambda \neq \lambda^*$ then $\mathrm{Res}^{\mathfrak{S}_n}_{\mathfrak{A}_n}(\chi_\lambda) = \mathrm{Res}^{\mathfrak{S}_n}_{\mathfrak{A}_n}(\chi_{\lambda^*})$ is irreducible. We will denote by ρ_λ this character of $\mathrm{Irr}(\mathfrak{A}_n)$. If λ is self-conjugate, then $\mathrm{Res}^{\mathfrak{S}_n}_{\mathfrak{A}_n}(\chi_\lambda)$ splits into two distinct constituents, denoted by $\rho_{\lambda,+}$ and $\rho_{\lambda,-}$. It follows immediately from Clifford's theory [2, III.2.14] that

$$\mathrm{Ind}^{\mathfrak{S}_n}_{\mathfrak{A}_n}(\rho_\lambda) = \chi_\lambda + \chi_{\lambda^*} \quad \text{and} \quad \mathrm{Ind}^{\mathfrak{S}_n}_{\mathfrak{A}_n}(\rho_{\lambda,+}) = \mathrm{Ind}^{\mathfrak{S}_n}_{\mathfrak{A}_n}(\rho_{\lambda,+}) = \chi_\lambda.$$

Recall that the conjugacy classes of $\mathfrak{S}_n$ are parametrized by the partitions of n (the parts of a partition λ of n give the cycle structure of the corresponding conjugacy class of $\mathfrak{S}_n$). If λ is a self-conjugate partition of n, then the class of $\mathfrak{S}_n$ labelled by the partition $\overline{\lambda}$ is contained in $\mathfrak{A}_n$, and splits into two classes, $\overline{\lambda}_+$ and $\overline{\lambda}_-$, of $\mathfrak{A}_n$. Moreover if c denotes a class of $\mathfrak{A}_n$ distinct from $\overline{\lambda}_+$ and $\overline{\lambda}_-$, then for $x \in c$, we have (cf [11, 2.5.13])

$$\rho_{\lambda,\pm}(x)=\chi_\lambda(x)/2,\quad \rho_{\lambda,\pm}(\overline{\lambda}_+)=x_\lambda\pm y_\lambda\quad\text{and}\quad \rho_{\lambda,\pm}(\overline{\lambda}_-)=x_\lambda\mp y_\lambda,$$

with $x_\lambda=\pm 1/2$ and y_λ is some non-zero complex number.

Remark 5.1. *Note that if x is a representative of a class of $\mathfrak{A}_n$ distinct from $\overline{\lambda}_+$ and $\overline{\lambda}_-$, then $\rho_{\lambda,\pm}(x)$ is a rational number, because $\rho_{\lambda,\pm}(x)=\chi_\lambda(x)/2$ with $\chi_\lambda(x)\in\mathbb{Z}$. However, $\rho_{\lambda,\pm}(x)$ is an algebraic integer. Hence it follows that $\rho_{\lambda,\pm}(x)\in\mathbb{Z}$.*

We can now prove:

Theorem 5.2. *Let n be a positive integer and p be an odd prime. Then $\mathfrak{A}_n$ has a p-basic set.*

Proof. Let $\mathcal{B}_\emptyset$ be the p-basic set of $\mathfrak{S}_n$ obtained in Theorem 1.1. We will prove that $\mathcal{B}_{\emptyset,\mathfrak{A}_n}$, the restriction of $\mathcal{B}_\emptyset$ to $\mathfrak{A}_n$, is a p-basic set of $\mathfrak{A}_n$. We denote by A the set of self-conjugate partitions λ of n such that $\overline{\lambda}$ is p-regular, and by S the set of sets $\{\lambda,\lambda^*\}$ with $\lambda\neq\lambda^*$ and $\chi_\lambda\in B$. For $x\in S$ we set $x=\{\lambda_x,\lambda_x^*\}$. By Lemma 3.4, if $\lambda=\lambda^*$, then $\chi_\lambda\in B$ if and only if $\lambda\in A$. We thus have

$$\mathcal{B}_\emptyset=\{\chi_\lambda\mid\lambda\in A\}\cup\{\chi_{\lambda_x},\chi_{\lambda_x^*}\mid x\in S\}.$$

It therefore follows that

$$\mathcal{B}_{\emptyset,\mathfrak{A}_n}=\{\rho_{\lambda,\pm}\mid\lambda\in A\}\cup\{\rho_{\lambda_x}\mid x\in S\}.$$

We will now prove that the family $\mathcal{B}_{\emptyset,\mathfrak{A}_n}^{p\text{-reg}}$ is free. Let $a_{\lambda,\pm}$ $(\lambda\in A)$, a_x $(x\in S)$ be integers such that

$$\sum_{\lambda\in A}(a_{\lambda,+}\rho_{\lambda,+}^{p\text{-reg}}+a_{\lambda,-}\rho_{\lambda,-}^{p\text{-reg}})+\sum_{x\in S}a_x\rho_{\lambda_x}^{p\text{-reg}}=0. \tag{3}$$

For every class function φ of $\mathfrak{A}_n$, we have $\operatorname{Ind}_{\mathfrak{A}_n}^{\mathfrak{S}_n}(\varphi^{p\text{-reg}})=(\operatorname{Ind}_{\mathfrak{A}_n}^{\mathfrak{S}_n}(\varphi))^{p\text{-reg}}$. Hence, inducing to $\mathfrak{S}_n$ Relation (3), we obtain

$$\sum_{\lambda\in A}(a_{\lambda,+}+a_{\lambda,-})\chi_\lambda^{p\text{-reg}}+\sum_{x\in S}a_x(\chi_{\lambda_x}^{p\text{-reg}}+\chi_{\lambda_x^*}^{p\text{-reg}})=0.$$

Using the fact that $\mathcal{B}_\emptyset^{p\text{-reg}}$ is free, we deduce that $a_x=0$ for all $x\in S$ and $a_{\lambda,+}=-a_{\lambda,-}$ for all $\lambda\in A$. Therefore, Relation (3) gives

$$\sum_{\lambda\in A}a_{\lambda,+}(\rho_{\lambda,+}^{p\text{-reg}}-\rho_{\lambda,-}^{p\text{-reg}})=0.$$

We now fix $\lambda_0\in A$. The partition $\overline{\lambda}_0$ is p-regular, so that the class of $\mathfrak{S}_n$ corresponding to $\overline{\lambda}_0$ is p-regular. It follows that the classes $\overline{\lambda}_{0,\pm}$ are also p-regular. For $\lambda\in A$ distinct from λ_0 we have $\rho_{\lambda,+}^{p\text{-reg}}(\overline{\lambda}_{0,+})=\rho_{\lambda,-}^{p\text{-reg}}(\overline{\lambda}_{0,+})$. Moreover $\rho_{\lambda_0,\pm}^{p\text{-reg}}(\overline{\lambda}_{0,+})=x_{\lambda_0}\pm y_{\lambda_0}$. It follows that

$$2a_{\lambda,+}y_{\lambda_0}=0.$$

But $y_{\lambda_0}\neq 0$ implies that $a_{\lambda,+}=0$, so that $a_{\lambda,+}=-a_{\lambda,-}=0$ for all $\lambda\in A$. Hence $\mathcal{B}_{\emptyset,\mathfrak{A}_n}^{p\text{-reg}}$ is free.

We will now prove that the family $\mathcal{B}_{\emptyset,\mathfrak{A}_n}^{p\text{-reg}}$ generates $\operatorname{Irr}(\mathfrak{A}_n)^{p\text{-reg}}$ over $\mathbb{Z}$. Let μ be a partition of n. Since $\mathcal{B}_\emptyset$ is a p-basic set, we have

$$\chi_\mu^{p\text{-reg}}=\sum_{\lambda\in A}a_{\mu,\lambda}\chi_\lambda^{p\text{-reg}}+\sum_{x\in S}(a_{\mu,x}\chi_{\lambda_x}^{p\text{-reg}}+b_{\mu,x}\chi_{\lambda_x^*}^{p\text{-reg}}), \tag{4}$$

for some integers $a_{\mu,\lambda}$ $(\lambda\in A)$, $a_{\mu,x}$ and $b_{\mu x}$ $(x\in S)$.

For every class function φ of $\mathfrak{S}_n$, we have $\operatorname{Res}_{\mathfrak{A}_n}^{\mathfrak{S}_n}(\varphi^{p\text{-reg}}) = (\operatorname{Res}_{\mathfrak{A}_n}^{\mathfrak{S}_n}(\varphi))^{p\text{-reg}}$.

We first suppose that $\mu \neq \mu^*$. Hence restricting to $\mathfrak{A}_n$ Relation (4), we obtain

$$\rho_\mu^{p\text{-reg}} = \sum_{\lambda \in A} a_{\mu,\lambda}(\rho_{\lambda,+}^{p\text{-reg}} + \rho_{\lambda,-}^{p\text{-reg}}) + \sum_{x \in S}(a_{\mu,x} + b_{\mu,x})\rho_{\lambda_x}^{p\text{-reg}},$$

so that $\rho_\mu^{p\text{-reg}}$ is a $\mathbb{Z}$-linear combination of elements of $\mathcal{B}_{\emptyset,\mathfrak{A}_n}^{p\text{-reg}}$.

We now suppose that μ is a self-conjugate partition not lying in A. In Relation (4), we have $a_{\mu,x} = b_{\mu,x}$ because χ_μ is ε-stable. Moreover $\overline{\mu}$ labels a p-singular class of $\mathfrak{S}_n$. Thus $\rho_{\mu,+}^{p\text{-reg}} = \rho_{\mu,-}^{p\text{-reg}}$, and it follows that

$$2\rho_\mu^{p\text{-reg}} = \sum_{\lambda \in A} a_{\mu,\lambda}(\rho_{\lambda,+}^{p\text{-reg}} + \rho_{\lambda,-}^{p\text{-reg}}) + \sum_{x \in S} 2a_{\mu,x}\rho_{\lambda_x}^{p\text{-reg}}.$$

Let $\lambda_0 \in A$. Evaluating the preceding relation on the class $\overline{\lambda}_{0,+}$, we obtain

$$2x_{\lambda_0}a_{\mu,\lambda_0} = 2\rho_\mu^{p\text{-reg}}(\overline{\lambda}_{0,+}) - \sum_{\lambda \neq \lambda_0} 2a_{\mu,\lambda}\,\rho_{\lambda,+}^{p\text{-reg}}(\overline{\lambda}_{0,+}) - \sum_{x \in S} 2a_{\mu,x}\,\rho_{\lambda_x}^{p\text{-reg}}(\overline{\lambda}_{0,+}) \in 2\mathbb{Z},$$

because $\rho_\mu^{p\text{-reg}}(\overline{\lambda}_{0,+}) \in \mathbb{Z}$ and $\rho_{\lambda,+}^{p\text{-reg}}(\overline{\lambda}_{0,+}) \in \mathbb{Z}$ for $\lambda \neq \lambda_0$ by Remark 5.1. However, $2x_{\lambda_0} = \pm 1$, so that $a_{\mu,\lambda_0} = 2a'_{\mu,\lambda}$ with $a'_{\mu,\lambda} \in \mathbb{Z}$. We therefore deduce that

$$\rho_\mu^{p\text{-reg}} = \sum_{\lambda \in A} a'_{\mu,\lambda}(\rho_{\lambda,+}^{p\text{-reg}} + \rho_{\lambda,-}^{p\text{-reg}}) + \sum_{x \in S} a_{\mu,x}\rho_{\lambda_x}^{p\text{-reg}}.$$

Since the result is obvious if $\mu \in A$, the result follows. □

6. Consequences for the decomposition matrices of $\mathfrak{A}_n$

Let G be a finite group and p a prime. We denote by $\operatorname{IBr}(G)$ the set of irreducible Brauer characters of G as in [2, IV.2]. Then, for every $\chi \in \operatorname{Irr}(G)$ and $\theta \in \operatorname{IBr}(G)$, there are non-negative integers $d_{\chi,\theta}$ (which are uniquely determined), such that

$$\chi^{p\text{-reg}} = \sum_{\theta \in \operatorname{IBr}(G)} d_{\chi,\theta}\,\theta.$$

The matrix $D = (d_{\chi,\theta})_{\chi \in \operatorname{Irr}(G), \theta \in \operatorname{IBr}(G)}$ is the so-called *p-decomposition matrix* of G [2, I.17]. In this section, we propose to deduce some results on the p-decomposition matrix of $\mathfrak{A}_n$ knowing that of $\mathfrak{S}_n$.

6.1. **Reduction of the problem.**

We keep the notation as above. Let B be a p-basic set of G. We set $D_B = (d_{\chi,\theta})_{\chi \in B, \theta \in \operatorname{IBr}(G)}$. Then D_B is a unimodular matrix. Moreover, D_B determines D, because we have

$$D = P_B D_B,$$

where, P_B is the matrix with coefficients $p_{\chi,\varphi}$ ($\chi \in \operatorname{Irr}(G)$, $\varphi \in B$), such that

$$\chi^{p\text{-reg}} = \sum_{\varphi \in B} p_{\chi,\varphi}\,\varphi^{p\text{-reg}}.$$

Note that, if $G = \mathfrak{S}_n$ and $B = \mathcal{B}_\emptyset$, then we know the coefficients $p_{\chi,\varphi}$ in the matrix P_B, at least theoretically. They come from their analogues $a_{\chi,\varphi}$ in $G_{p,w}$, described in the proof of Proposition 4.2.

Proposition 6.1. *Let p be an odd prime. Let B be an ε-stable p-basic set of $\mathfrak{S}_n$, where ε denotes the sign character of $\mathfrak{S}_n$. We denote by D_B the restriction of the p-decomposition matrix of $\mathfrak{S}_n$ to B as above. Then ε acts on $\operatorname{IBr}(\mathfrak{S}_n)$, the rows of D_B and the columns of D_B, and these three operations are equivalent.*

Proof. We denote by $\operatorname{Fix}_\varepsilon(B)$ (respectively $\operatorname{Fix}_\varepsilon(\operatorname{IBr}(\mathfrak{S}_n))$) the set of ε-stable characters in B (respectively in $\operatorname{IBr}(\mathfrak{S}_n)$). In order to prove the assumption, we have to prove

$$|\operatorname{Fix}_\varepsilon(B)| = |\operatorname{Fix}_\varepsilon(\operatorname{IBr}(\mathfrak{S}_n))|.$$

For ease of notation, we will, throughout the proof, write $\hat{\varphi}$ for $\varphi^{p\text{-reg}}$ ($\varphi \in \operatorname{Irr}(\mathfrak{S}_n)$).

We set $D_B = (d_{\chi,\theta})_{\chi\in B,\theta\in\operatorname{IBr}(\mathfrak{S}_n)}$ and $D_{B,\varepsilon} = (d_{\varepsilon\chi,\hat{\varepsilon}\theta})_{\varepsilon\chi\in B,\hat{\varepsilon}\theta\in\operatorname{IBr}(\mathfrak{S}_n)}$. We have

$$\widehat{\varepsilon\chi} = \sum_{\theta\in\operatorname{IBr}(\mathfrak{S}_n)} d_{\varepsilon\chi,\theta}\theta.$$

Hence, it follows that

$$\begin{aligned}\hat{\chi} &= \sum_{\theta\in\operatorname{IBr}(\mathfrak{S}_n)} d_{\varepsilon\chi,\theta}\hat{\varepsilon}\theta \\ &= \sum_{\hat{\varepsilon}\theta'\in\operatorname{IBr}(\mathfrak{S}_n)} d_{\varepsilon\chi,\hat{\varepsilon}\theta'}\theta' \\ &= \sum_{\theta'\in\operatorname{IBr}(\mathfrak{S}_n)} d_{\varepsilon\chi,\hat{\varepsilon}\theta'}\theta'.\end{aligned}$$

Since $\operatorname{IBr}(\mathfrak{S}_n)$ is a $\mathbb{Z}$-basis of $\mathbb{Z}\operatorname{IBr}(\mathfrak{S}_n)$, we deduce that $d_{\chi,\theta} = d_{\varepsilon\chi,\hat{\varepsilon}\theta}$, that is

$$D_B = D_{B,\varepsilon}.$$

Furthermore, $D_{B,\varepsilon}$ is obtained from D_B by a permutation of the rows and of the columns. Thus, there are permutation matrices P and Q such that

$$D_{B,\varepsilon} = PD_BQ.$$

We deduce that $D_B = PD_BQ$. Furthermore, we have $|B| = |\operatorname{IBr}(\mathfrak{S}_n)| = m$. Since B is a $\mathbb{Z}$-basis of $\mathbb{Z}\operatorname{IBr}(\mathfrak{S}_n)$, we have $D_B \in \operatorname{GL}_m(\mathbb{Z})$. Thus, we have

$$Q^{-1} = D_B^{-1}PD_B.$$

It follows that $\operatorname{Tr}(Q^{-1}) = \operatorname{Tr}(P)$. Since P, Q, P^{-1} and Q^{-1} are permutation matrices, we deduce that they have the same trace. Furthermore, the number of fixed-point of the permutation induced by P is the number of fixed rows. Hence, we have $\operatorname{Tr}(P) = |\operatorname{Fix}_\varepsilon(B)|$. Moreover, we have $D_B^{-1} = QD_B^{-1}P$. We then deduce that $\operatorname{Tr}(Q) = |\operatorname{Fix}_\varepsilon(\operatorname{IBr}(\mathfrak{S}_n))|$, and the result follows. ☐

We denote by $\mathcal{B}_\emptyset$ the ε-stable p-basic set of $\mathfrak{S}_n$ obtained in Theorem 1.1 and by $\mathcal{B}_{\mathfrak{A}_n,\emptyset}$ the restriction of $B_\emptyset$ to $\mathfrak{A}_n$. As above, if λ is a partition of n, we denote by α_λ its p-quotient. As above, we set

$$\Lambda_\emptyset = \{\lambda \vdash n \mid \alpha_\lambda^{(p-1)/2} = \emptyset\}.$$

We denote by $\Lambda_{\emptyset,1}$ the set of $\lambda \in \Lambda_\emptyset$ such that $\lambda = \lambda^*$ and by $\Lambda_{\emptyset,2}$ the set of sets $\{\lambda, \lambda^*\}$ for $\lambda \in \Lambda_\emptyset$ with $\lambda \neq \lambda^*$. For $w \in \Lambda_{\emptyset,2}$, we fix $\lambda_w \in w$. In particular, we have $w = \{\lambda_w, \lambda_w^*\}$. In Theorem 1.1 and in the proof of Theorem 5.2, we proved that

$$\mathcal{B}_\emptyset = \{\chi_\lambda \mid \lambda \in \Lambda_\emptyset\} \quad \text{and} \quad \mathcal{B}_{\mathfrak{A}_n,\emptyset} = \{\rho_{\lambda,\pm} \mid \lambda \in \Lambda_{\emptyset,1}\} \cup \{\rho_{\lambda_w} \mid w \in \Lambda_{\emptyset,2}\},$$

where $\operatorname{Res}_{\mathfrak{A}_n}^{\mathfrak{S}_n}(\chi_\lambda) = \rho_{\lambda,+} + \rho_{\lambda,-}$ for $\lambda \in \Lambda_{\emptyset,1}$, and $\rho_{\lambda_w} = \operatorname{Res}_{\mathfrak{A}_n}^{\mathfrak{S}_n}(\chi_{\lambda_w})$ for $w \in \Lambda_{\emptyset,2}$.

By Proposition 6.1, we can take $\Phi : \mathcal{B}_\emptyset \to \operatorname{IBr}(\mathfrak{S}_n)$ an ε-equivariant bijection. For λ, $\mu \in \Lambda_\emptyset$, we set $\theta_\mu = \Phi(\chi_\mu)$ and $d_{\lambda,\mu} = d_{\chi_\lambda,\theta_\mu}$. By Clifford's modular theory [10, 9.18] (which could be applied because the quotient $\mathfrak{S}_n/\mathfrak{A}_n$ is cyclic of order prime to p) we know how the θ_μ's restrict from $\mathfrak{S}_n$ to $\mathfrak{A}_n$. If $\mu \in \Lambda_{\emptyset,1}$, then $\operatorname{Res}^{\mathfrak{S}_n}_{\mathfrak{A}_n}(\theta_\mu)$ is the sum of two irreducible Brauer characters of $\mathfrak{A}_n$, which we denote by $\theta'_{\mu,+}$ and $\theta'_{\mu,-}$. If $w \in \Lambda_{\emptyset,2}$, then $\theta'_{\mu_w} = \operatorname{Res}^{\mathfrak{S}_n}_{\mathfrak{A}_n}(\theta_{\mu_w}) \in \operatorname{IBr}(\mathfrak{A}_n)$. We then have

$$\operatorname{IBr}(\mathfrak{A}_n) = \{\theta'_{\mu,\pm} \mid \mu \in \Lambda_{\emptyset,1}\} \cup \{\theta'_{\mu_w} \mid w \in \Lambda_{\emptyset,2}\}.$$

For λ, $\mu \in \Lambda_{\emptyset,1}$ and w, $w' \in \Lambda_{\emptyset,2}$, there are non-negative integers $d'_{\lambda;\pm,\mu_w}$, $d'_{\lambda;\pm,\mu,\pm}$, $d'_{\lambda_{w'},\mu_w}$ and $d'_{\lambda_{w'},\mu;\pm}$ such that

$$\rho^{p\text{-reg}}_{\lambda,\pm} = \sum_{\mu\in\Lambda_{\emptyset,1}} (d'_{\lambda;\pm,\mu;+}\theta'_{\mu,+} + d'_{\lambda;\pm,\mu;-}\theta'_{\mu,-}) + \sum_{w\in\Lambda_{\emptyset,2}} d'_{\lambda;\pm,\mu_w}\theta_{\mu_w},$$

$$\rho^{p\text{-reg}}_{\lambda_{w'}} = \sum_{\mu\in\Lambda_{\emptyset,1}} (d'_{\lambda_{w'},\mu;+}\theta'_{\mu,+} + d'_{\lambda_{w'},\mu;-}\theta'_{\mu,-}) + \sum_{w\in\Lambda_{\emptyset,2}} d'_{\lambda_{w'},\mu_w}\theta_{\mu_w}.$$

We then obtain the following about the p-decomposition numbers of $\mathfrak{S}_n$ and $\mathfrak{A}_n$:

Theorem 6.2. *With the above notations, we have, for all λ, $\mu \in \Lambda_{\emptyset,1}$ and for all w, $w' \in \Lambda_{\emptyset,2}$,*
(i)

$$d_{\lambda,\mu_w} = d_{\lambda,\mu^*_w}$$
$$d_{\lambda_{w'},\mu} = d_{\lambda^*_{w'},\mu}$$
$$d_{\lambda_{w'},\mu_w} = d_{\lambda^*_{w'},\mu^*_w} \quad \textit{and} \quad d_{\lambda_{w'},\mu^*_w} = d_{\lambda^*_{w'},\mu_w},$$

(ii)

$$d'_{\lambda;+,\mu;+} + d'_{\lambda;-,\mu;+} = d_{\lambda,\mu} = d'_{\lambda;+,\mu;-} + d'_{\lambda;-,\mu;-}$$
$$d'_{\lambda;+,\mu;+} = d'_{\lambda;-,\mu;-} \quad \textit{and} \quad d'_{\lambda;+,\mu;-} = d'_{\lambda;-,\mu;+}$$
$$d'_{\lambda;+,\mu_w} = d'_{\lambda;-,\mu_w} = d_{\lambda,\mu_w},$$

(iii)

$$d_{\lambda_{w'},\mu_w} + d_{\lambda_{w'},\mu^*_w} = d'_{\lambda_{w'},\mu_w}$$
$$d'_{\lambda_{w'},\mu;+} = d'_{\lambda_{w'},\mu;-} = d_{\lambda_{w'},\mu}.$$

Proof. (i) This is a direct consequence of our parametrizations and our choice of Φ.
(ii) Since $\operatorname{Res}^{\mathfrak{S}_n}_{\mathfrak{A}_n}(\chi_\lambda) = \rho_{\lambda,+} + \rho_{\lambda,-}$, $\operatorname{Res}^{\mathfrak{S}_n}_{\mathfrak{A}_n}(\theta_\mu) = \theta'_{\mu,+} + \theta'_{\mu,-}$ and $\operatorname{Res}^{\mathfrak{S}_n}_{\mathfrak{A}_n}(\theta_{\mu_w}) = \operatorname{Res}^{\mathfrak{S}_n}_{\mathfrak{A}_n}(\theta_{\mu^*_w}) = \theta'_{\mu_w}$, we immediately get

$d'_{\lambda;+,\mu;+} + d'_{\lambda;-,\mu;+} = d_{\lambda,\mu} = d'_{\lambda;+,\mu;-} + d'_{\lambda;-,\mu;-}$ and $d_{\lambda,\mu_w} + d_{\lambda,\mu^*_w} = d'_{\lambda;+,\mu_w} + d'_{\lambda;-,\mu_w}$.

Now, if we write τ for the automorphism of $\mathfrak{A}_n$ induced by conjugation by a transposition of $\mathfrak{S}_n$, then τ also acts on $\operatorname{Irr}(\mathfrak{A}_n)$ and $\operatorname{IBr}(\mathfrak{A}_n)$, and we have

$$\tau(\theta'_{\mu,+}) = \theta'_{\mu,-}, \quad \tau(\theta'_{\mu_w}) = \theta'_{\mu_w} \quad \text{and} \quad \tau(\rho_{\lambda,+}) = \rho_{\lambda,-}.$$

This implies

$$d'_{\lambda;+,\mu;+} = d'_{\lambda;-,\mu;-}, \quad d'_{\lambda;+,\mu;-} = d'_{\lambda;-,\mu;+} \quad \text{and} \quad d'_{\lambda;+,\mu_w} = d'_{\lambda;-,\mu_w}.$$

Together with (i) and the above, this finally gives $d'_{\lambda;+,\mu_w} = d'_{\lambda;-,\mu_w} = d_{\lambda,\mu_w}$.
(iii) Since $\operatorname{Res}^{\mathfrak{S}_n}_{\mathfrak{A}_n}(\chi_{\lambda_{w'}}) = \operatorname{Res}^{\mathfrak{S}_n}_{\mathfrak{A}_n}(\chi_{\lambda^*_{w'}}) = \rho_{\lambda_{w'}}$, $\operatorname{Res}^{\mathfrak{S}_n}_{\mathfrak{A}_n}(\theta_\mu) = \theta'_{\mu,+} + \theta'_{\mu,-}$ and $\operatorname{Res}^{\mathfrak{S}_n}_{\mathfrak{A}_n}(\theta_{\mu_w}) = \operatorname{Res}^{\mathfrak{S}_n}_{\mathfrak{A}_n}(\theta_{\mu^*_w}) = \theta'_{\mu_w}$, the desired equalities follow easily.

□

We now show on a picture how to use Theorem 6.2 to obtain decomposition numbers of $\mathfrak{A}_n$, knowing those of $\mathfrak{S}_n$ (or conversely). We denote by $D_{\mathcal{B}_\emptyset}$ the restriction to the basic set $\mathcal{B}_\emptyset$ of the p-decomposition matrix of $\mathfrak{S}_n$, and $D_{\mathcal{B}_{\emptyset,\mathfrak{A}_n}}$ the restriction to $\mathcal{B}_{\emptyset,\mathfrak{A}_n}$ of the decomposition matrix of $\mathfrak{A}_n$. Using the notations of Theorem 6.2, we write $a = d_{\lambda,\mu}$, $b = d_{\lambda,\mu_w}$, $c = d_{\lambda_{w'},\mu}$, $d = d_{\lambda_{w'},\mu_w}$, $e = d_{\lambda_{w'},\mu_w^*}$, $\alpha = d'_{\lambda;+,\mu;+}$ and $\beta = d'_{\lambda;+,\mu;-}$. We also write $a_1 = |\Lambda_{\emptyset,1}|$ and $a_2 = |\Lambda_{\emptyset,2}|$. We start, at the level of $\mathfrak{S}_n$, with



Then, by Theorem 6.2, we obtain, at the level of $\mathfrak{A}_n$,



Note that, by Theorem 6.2, we have $\alpha + \beta = a$.

6.2. **Consequences.** In this section, we suppose that we know the decomposition matrix D of $\mathfrak{S}_n$. We denote by $D_{\mathcal{B}_\emptyset}$ the restriction of D to the p-basic set $\mathcal{B}_\emptyset$ as above. Let $D_{n,p}$ be the submatrix of $D_{\mathcal{B}_\emptyset}$ whose entries are all the entries of $D_{\mathcal{B}_\emptyset}$ which lie at the intersection of an ε-stable row and an ε-column of $D_{\mathcal{B}_\emptyset}$. We can restrict this matrix to $\mathfrak{A}_n$. We thus obtain a matrix $D'_{n,p}$ which has twice as many rows and columns as $D_{n,p}$.

Theorem 6.3. *We keep the notation as above. We suppose that the p-decomposition matrix D of $\mathfrak{S}_n$ and the matrix $D'_{n,p}$ are known. Then we can construct the p-decomposition matrix of $\mathfrak{A}_n$.*

Proof. As above, we denote by $D_{\mathcal{B}_{\emptyset,\mathfrak{A}_n}}$ the restriction of the p-decomposition matrix of $\mathfrak{A}_n$ to the p-basic set $\mathcal{B}_{\emptyset,\mathfrak{A}_n}$ (obtained in Theorem 5.2). As explained at the begining of Section 6, to compute the p-decomposition matrix of $\mathfrak{A}_n$, it is sufficient to compute the matrix $D_{\mathcal{B}_{\emptyset,\mathfrak{A}_n}}$ (again, provided we know the matrix $P_{\mathcal{B}_{\emptyset,\mathfrak{A}_n}}$). Furthermore, using Theorem 6.2, every coefficient of $D_{\mathcal{B}_{\emptyset,\mathfrak{A}_n}}$ which does not lie in $D'_{n,p}$ can be obtained from D. The result follows. □

The above theorem shows that, in order to deduce the p-decomposition matrix of $\mathfrak{A}_n$ from that of $\mathfrak{S}_n$, we just have to understand the matrix $D'_{n,p}$, which is a small matrix compared to the p-decomposition matrix of $\mathfrak{A}_n$.

Following [11, §6.3], we say that a matrix has wedge shape if its rows and its columns can be ordered in such a way that the resulting matrix is a lower triangular matrix with diagonal entries equal to 1.

Lemma 6.4. *We suppose that $\mathfrak{S}_n$ has a p-basic set B' satisfying the properties of Theorem 1.1, and such that the restriction matrix $D_{B'}$ has wedge shape. Then the matrix $D_{n,p}$ defined above (and obtained from the p-basic set $\mathcal{B}_\emptyset$ constructed in Theorem 1.1) has wedge shape.*

Proof. Let $\chi_\lambda \in B'$ be such that $\lambda = \lambda^*$. Since B' has Property (2) of Theorem 1.1, it follows that $\overline{\lambda}$ is p-regular. Furthermore, by Lemma 3.4, we see that $\lambda \in \Lambda_\emptyset$. Hence, the ε-stable characters in B' are the same as those in $\mathcal{B}_\emptyset$. The result follows. □

Remark 6.5. *The p-basic set $\mathcal{B}_\emptyset$ does not have wedge shape. The first counter-example is given by $\mathfrak{S}_6$ with $p = 3$. In this case, we have*

$$D_{\mathcal{B}_\emptyset} = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 0 & 0 \end{bmatrix},$$

which has no wedge shape. However, there is, at the moment, no example where Lemma 6.4 does not apply. Indeed, one can check, using GAP *[8], that, for $1 \leq n \leq 18$ and for any odd prime, there always exists a p-basic set B' as in Lemma 6.4. We can then conjecture that such a p-basic set of $\mathfrak{S}_n$ always exists and that $D_{n,p}$ has wedge shape.*

Lemma 6.6. *With the above notation, if $D_{n,p}$ has wedge shape, then so does $D'_{n,p}$.*

Proof. We suppose that $D_{n,p}$ is a matrix of size r and has wedge shape. Hence we can order the rows and columns in such a way that the resulting matrix is lower triangular with diagonal entries equal to 1. This induces permutations τ_r and τ_c on the rows and the columns of $D_{n,p}$ respectively. By definition, $D'_{n,p}$ is a matrix of size $2r$. Now, we first regroup the rows and the columns of $D'_{n,p}$ in corresponding pairs, according to Theorem 6.2, and label the pairs by $(2k-1,2k)$ for $1 \leq k \leq r$. We then apply the permutations of the rows and the columns defined by $(2k-1,2k) \mapsto (2\tau_r(k)-1, 2\tau_r(k))$ and $(2k-1,2k) \mapsto (2\tau_c(k)-1, 2\tau_c(k))$ respectively. The resulting matrix is thus lower triangular by blocks, and the blocks on the diagonal have size 2. To every coefficient $d_{i,j}$ of $D_{n,p}$, we associate the submatrix $D_{i,j}$ of $D'_{n,p} = (d'_{i,j})$ given by

$$D_{i,j} = \begin{bmatrix} d'_{2i-1,2j-1} & d'_{2i-1,2j} \\ d'_{2i,2j-1} & d'_{2i,2j} \end{bmatrix}.$$

In particular, the 2 by 2 blocks on the diagonal of $D'_{n,p}$ are the $D_{i,i}$'s, $1 \leq i \leq r$. Moreover, Theorem 6.2 asserts that $d'_{2i-1,2j-1} = d'_{2i,2j}$, $d'_{2i-1,2j} = d'_{2i,2j-1}$ and $d'_{2i-1,2j-1} + d'_{2i-1,2j} = d_{i,j}$. Since $d_{i,i} = 1$, we thus deduce that

$$D_{i,i} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \quad \text{or} \quad D_{i,i} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

We can now re-order the rows of $D'_{n,p}$ as follows: for every $1 \leq i \leq r$, if $D_{i,i}$ is not the identity matrix, then we permute the rows $2i-1$ and $2i$ of $D'_{n,p}$. The resulting matrix has wedge shape, as required. ☐

Remark 6.7. *Note that, if $D_{n,p}$ has wedge shape, then, using our p-basic set $\mathcal{B}_{\emptyset,\mathfrak{A}_n}$ obtained in Theorem 5.2, we have a way to parametrize the simple module of $\mathfrak{A}_n$ in characteristic p which are not the restriction of a simple module of $\mathfrak{S}_n$.*

Remark 6.8. *We now discuss some problems and questions. We suppose that the matrix $D_{n,p}$ is known and has size r. Let $d_{i,j}$ be the (i,j)-coefficient of $D_{n,p}$. As in the proof of Lemma 6.6, we can associate to this coefficient the 2 by 2 matrix $D_{i,j}$. Following Theorem 6.3, to construct the p-decomposition matrix of $\mathfrak{A}_n$, it is sufficient to compute all the matrices $D_{i,j}$ $(1 \leq i,j \leq r)$. Furthermore, by Theorem 6.2, we have*

$$D_{i,j} = \begin{bmatrix} a & b \\ b & a \end{bmatrix},$$

with $a+b = d_{i,j}$. Hence, the computation of the p-decomposition matrix of $\mathfrak{A}_n$ is reduced to the following problems.

- *For $1 \leq i,j \leq r$, we have to determine the integers a and b appearing in the matrix $D_{i,j}$.*
- *If the coefficients a and b appearing in all the matrices $D_{i,j}$ $(1 \leq i,j \leq r)$ are known, then a p-decomposition matrix of $\mathfrak{A}_n$ is known, up to some choices. We have to develop a rule to fix theses choices.*

Acknowledgements. Part of this work was done during a visit of the first author in Lausanne. The first author gratefully acknowledges financial support by the IGAT (EPFL). We wish to thank Michael Cuntz for many helpful and valuable discussions on the subject. Some of the ideas in this paper originated from discussions with him during a postdoctoral stay of the first author at the University of

Kaiserlautern. We also thank Meinolf Geck for his reading of the manuscript, and for his suggestions and helpful comments on this paper.

References

[1] O. Brunat. Basic sets in defining characteristic for general linear groups of small rank. *J. Pure Appl. Algebra*, doi:10.1016/j.jpaa.2008.09.006, 2008.

[2] W. Feit. *The representation theory of finite groups*, volume 25 of *North-Holland Mathematical Library*. North-Holland Publishing Co., Amsterdam, 1982.

[3] M. Geck. Irreducible Brauer characters of the 3-dimensional special unitary groups in non-defining characteristic. *Comm. Algebra*, 18(2):563–584, 1990.

[4] M. Geck. Basic sets of Brauer characters of finite groups of Lie type. II. *J. London Math. Soc. (2)*, 47(2):255–268, 1993.

[5] M. Geck. Basic sets of Brauer characters of finite groups of Lie type. III. *Manuscripta Math.*, 85(2):195–216, 1994.

[6] M. Geck and G. Hiss. Basic sets of Brauer characters of finite groups of Lie type. *J. Reine Angew. Math.*, 418:173–188, 1991.

[7] J.-B. Gramain. On defect groups for generalized blocks of the symmetric group. *J. London Math. Soc.*, 78(1):155–171, 2008.

[8] The Gap Group. Groups, algorithms, and programming, version 4.4.9,. *http://www.gap-system.org, 2006.*

[9] G. Hiss. On the decomposition numbers of $G_2(q)$. *J. Algebra*, 120(2):339–360, 1989.

[10] B. Huppert and N. Blackburn. *Finite groups. II*, volume 242 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1982. AMD, 44.

[11] G. James and A. Kerber. *The representation theory of the symmetric group*, volume 16 of *Encyclopedia of Mathematics and its Applications*. Addison-Wesley Publishing Co., Reading, Mass., 1981. With a foreword by P. M. Cohn, With an introduction by Gilbert de B. Robinson.

[12] B. Külshammer, J. Olsson, and G.R. Robinson. Generalized blocks for symmetric groups. *Invent. Math.*, 151(3):513–552, 2003.

[13] M. Osima. On the representations of the generalized symmetric group. *Math. J. Okayama Univ.*, 4:39–56, 1954.

[14] Götz Pfeiffer. Character tables of Weyl groups in GAP. *Bayreuth. Math. Schr.*, 47:165–222, 1994.

Ruhr-Universität Bochum, Fakultät für Mathematik, Raum NA 2/33, D-44780 Bochum,
E-mail address: `Olivier.Brunat@ruhr-uni-bochum.de`

École Polytechnique Fédérale de Lausanne, Institut de géométrie, algèbre et topologie, Bâtiment de Chimie (BCH), CH-1015 Lausanne,
E-mail address: `jean-baptiste.gramain@epfl.ch`